


CGP



GCSE Chemistry

For AQA (Grade 9-1)

New!

Exam Practice Answer Book

Contents

Topic 1 — Atomic Structure and the Periodic Table	2
Topic 2 — Bonding, Structure and Properties of Matter	4
Topic 3 — Quantitative Chemistry	7
Topic 4 — Chemical Changes	9
Topic 5 — Energy Changes	11
Topic 6 — The Rate and Extent of Chemical Change	11
Topic 7 — Organic Chemistry	13
Topic 8 — Chemical Analysis	15
Topic 9 — Chemistry of the Atmosphere	16
Topic 10 — Using Resources	17
Mixed Questions	18

Published by CGP

www.cgpbooks.co.uk

Printed by Elanders Ltd, Newcastle upon Tyne.

Clipart from Corel®

Text, design, layout and original illustrations © Coordination Group Publications Ltd. (CGP) 2016

All rights reserved.

Topic 1 — Atomic Structure and the Periodic Table

Page 1 — Atoms

Warm-up

The radius of an atom is approximately **0.1** nanometres.
The radius of the nucleus is around 1×10^{-14} metres. That's about **1/10 000** of the radius of an atom.

An atom doesn't have an overall **charge** as it has equal numbers of **protons/electrons** and **electrons/protons**.

1.1 nucleus [1 mark]

1.2 -1 [1 mark]

1.3 neutron: 0 charge [1 mark]

proton: +1 charge [1 mark]

2.1 mass number = 39 [1 mark]

2.2 atomic number = 19 [1 mark]

2.3 protons = 19 [1 mark]

neutrons = mass number - atomic number

= 39 - 19 = **20** [1 mark]

electrons = 19 [1 mark]

Page 2 — Elements

1.1 Atoms are the smallest part of an element that can exist [1 mark].

1.2 They have the same number of protons / 17 protons [1 mark] but a different number of neutrons / $^{35}_{17}\text{Cl}$ has 2 less neutrons than $^{37}_{17}\text{Cl}$ [1 mark].

2.1

Isotope	No. of Protons	No. of Neutrons	No. of Electrons
^{32}S	16	16	16
^{33}S	16	17	16
^{34}S	16	18	16
^{36}S	16	20	16

[3 marks — 1 mark for each correct column]

2.2 Relative atomic mass = $[(94.99 \times 32) + (0.75 \times 33) + (4.25 \times 34) + (0.01 \times 36)] \div (94.99 + 0.75 + 4.25 + 0.01)$
= $3209.29 \div 100 = 32.0929 = \mathbf{32.1}$ [2 marks for correct answer, otherwise one mark for using correct equation]

2.3 X and Z are isotopes [1 mark]. They have the same atomic number / same number of protons [1 mark] but different mass numbers / number of neutrons [1 mark].

Page 3 — Compounds

1.1 It contains two elements chemically combined [1 mark].

1.2 4 [1 mark]

A molecule of ammonia contains 1 nitrogen atom and 3 hydrogen atoms making a total of 4 atoms altogether.

2.1 sodium chloride [1 mark]

2.2 Any one of: **B.** NaCl / **C.** C_2H_4 / **E.** H_2O [1 mark]

It contains two or more elements chemically combined (in fixed proportions) [1 mark].

2.3 6 [1 mark]

C_2H_4 contains 2 carbon atoms and 4 hydrogen atoms.

2.4 Yes, a new compound has been made as the atoms in C_2H_6 are in different proportions to the atoms in C or F / there are a different number of hydrogen atoms in the molecule [1 mark].

Page 4 — Chemical Equations

Warm-up

1 True

2 False

3 True

4 True

1.1 sodium + chlorine → sodium chloride [1 mark]

1.2 $2\text{Na} + \text{Cl}_2 \rightarrow 2\text{NaCl}$ [1 mark]

2.1 $4\text{NH}_3 + 5\text{O}_2 \rightarrow 4\text{NO} + 6\text{H}_2\text{O}$ /

$2\text{NH}_3 + 2.5\text{O}_2 \rightarrow 2\text{NO} + 3\text{H}_2\text{O}$ [1 mark]

2.2 E.g. there are 7 oxygen atoms on the left hand side of the equation and only 6 on the right hand side [1 mark].

Page 5 — Mixtures and Chromatography

- 1.1 Mixture [1 mark]. Air consists of two or more elements or compounds [1 mark] that aren't chemically combined together [1 mark].
- 1.2 No [1 mark], as argon is an element in a mixture. Chemical properties are not affected by being in a mixture [1 mark].
- 2 How to grade your answer:
 Level 0: Nothing written worthy of credit [No marks].
 Level 1: Some explanation or description given but little detail and key information missing [1–2 marks].
 Level 2: Clear description of method and some explanation of results but some detail missing [3–4 marks].
 Level 3: A clear and detailed description of method and a full explanation of results [5–6 marks].
- Here are some points your answer may include:
Setting up the experiment
 Draw a line in pencil near the bottom of a piece of chromatography paper.
 Place a small sample of each ink on the pencil line.
 Pour a shallow layer of water / solvent into a beaker.
 Place the chromatography paper in the container.
 The water should be below the pencil line and the ink spots.
 Place a lid on the container and wait for the solvent to rise to near the top of the paper.
 Remove the paper from the container when the solvent has risen close to the top of the paper.
- Explanation of results
 A shows one spot, so only contains one dye.
 B shows two spots that have separated, so contains two dyes.
 C shows three spots that have separated, so contains three dyes.
 B and C are mixtures as they contain more than one element or compound not chemically combined together.
 B and C contain at least one of the same dyes

Page 6 — More Separation Techniques

- 1.1 Add water to the mixture to dissolve the potassium chloride [1 mark]. Filter the mixture. The chalk will stay on the filter paper, [1 mark] the dissolved potassium chloride will pass through [1 mark].
- 1.2 E.g. evaporate the potassium chloride solution to a much smaller volume and then leave it to cool [1 mark].
- 2.1 Add the mixture to methylbenzene. The sulfur will dissolve (the iron will not dissolve) [1 mark]. Filter the solution to obtain the insoluble iron [1 mark]. Evaporate the methylbenzene to obtain crystals of sulfur [1 mark].
- 2.2 No, the student is incorrect [1 mark]. The iron and sulfur are chemically combined in iron(II) sulfide / iron(II) sulfide is a compound [1 mark] so chemical methods would be needed to separate them out [1 mark].

Page 7 — Distillation

- 1 Simple distillation [1 mark]
- 2.1 Place a stopper / stopper with a thermometer in the top of the distillation flask [1 mark].
- 2.2 The solution is heated/boiled and the aspirin evaporates first as it has a lower boiling point than the impurity [1 mark]. There is cold water flowing through the (Liebig) condenser [1 mark]. This condenses the gaseous aspirin back into a liquid which is then collected [1 mark].
- 2.3 The aspirin has a boiling point greater than 100 °C / greater than the boiling point of water [1 mark]. So it would not evaporate [1 mark].

Pages 8-9 — The History of The Atom

Warm-up

New experimental evidence can disprove models — **True**
 Scientific models can be based on existing theories and new experimental evidence — **True**

Older scientific theories must be ignored when new ones are adopted — **False**

- 1.1 Tiny solid spheres that can't be divided [1 mark].
- 1.2 Plum pudding model — A positively charged 'ball' with negatively charged electrons in it [1 mark].
 Bohr's model — Electrons in fixed orbits surrounding a small positively charged nucleus [1 mark].
 Rutherford's nuclear model — A small positively charged nucleus surrounded by a 'cloud' of negative electrons [1 mark].
- 1.3 neutron [1 mark]
- 2.1 Most of the atom is "empty" space [1 mark].
- 2.2 Niels Bohr [1 mark]
- 3.1 Atoms are neutral / have no overall charge [1 mark].
 Therefore there must have been positive charge to balance the negative charge of the electrons [1 mark].
- 3.2 How to grade your answer:
 Level 0: Nothing written worthy of credit [No marks].
 Level 1: A brief description of either the nuclear or the 'plum pudding' model is given [1 to 2 marks].
 Level 2: A description of both the nuclear model and the plum pudding model is given and some comparisons made [3 to 4 marks].
 Level 3: A full comparison of the models is given and similarities and differences are clearly explained [5 to 6 marks].

Here are some points your answer may include:

Similarities

They both have areas of positive charge.

They both have electrons.

They are both neutral overall.

Differences

Positive charge isn't divided into protons in plum pudding model.

Plum pudding model does not have a nucleus but has a 'ball' of positive charge instead.

Plum pudding model does not have neutrons or protons, it only has electrons surrounded by a positive charge.

Plum pudding model does not have shells of electrons (surrounding nucleus), the electrons are arranged randomly within a sphere of positive charge.


Modern nuclear model has most of the mass concentrated in the nucleus but the plum pudding model has the mass spread evenly throughout the entire atom.

Page 10 — Electronic Structure

- 1.1 2,8,8,2 [1 mark]
- 1.2 The electrons in an atom occupy the lowest energy levels/ innermost shell first [1 mark]. The innermost shell/lowest energy level can hold 2 electrons [1 mark].

2.1 Chlorine: 2,8,7 [1 mark]

2.2



[1 mark for correct number of electrons, 1 mark for correct arrangement]

You don't have to have the electrons paired up on the diagram. As long as there is the same number of electrons on the same shells you get the marks.

2.3 Phosphorus/P [1 mark]

Page 11 — Development of The Periodic Table

- 1.1 He left gaps so that elements with similar properties were in the same group / for elements that had not yet been discovered [1 mark].
- 1.2 D. Between 2.4 and 7.2 g/cm³ [1 mark]. E. EkO₂ [1 mark]
 F. EkCl₄ [1 mark] G. Very slow [1 mark].
- 2.1 Protons (neutrons and electrons) had not been discovered / atomic numbers weren't known [1 mark].

- 2.2 Ar and K / Te and I [1 mark].
 2.3 Isotopes of an element have different numbers of neutrons / different atomic masses [1 mark], but the same chemical properties [1 mark].

Page 12 — The Modern Periodic Table

- 1.1 By atomic number / proton number [1 mark].
 1.2 Similar properties occur at regular intervals / there are repeating patterns in the properties of the elements [1 mark].
 1.3 They have the same number of outer shell electrons [1 mark].
 2.1 Group 2 [1 mark]. The atom has 2 outer shell electrons. [1 mark].
 2.2 Period 3 [1 mark]. The atom has 3 shells of electrons [1 mark].
 2.3 Magnesium/Mg [1 mark]
 2.4 Choose one from: beryllium / calcium / strontium / barium / radium [1 mark]

Page 13 — Metals and Non-Metals

- 1.1 Metals: Towards the left and bottom.
 Non-metals: Towards the right and top [1 mark].
 1.2 Elements that react to form positive ions are metals [1 mark].
 1.3 Any one from: e.g. good electrical conductor / good thermal conductor / strong / high boiling point / high melting point / malleable [1 mark].
 1.4 Both are metals that lose their (2 or 3) outer shell electrons [1 mark] to form positive ions [1 mark].
 2.1 Iron is a metal and therefore strong / good at conducting heat/electricity / has a high boiling/melting point [1 mark for each property up to a maximum of 2 marks]. Sulfur is a non-metal and therefore is more brittle / has a lower density / doesn't conduct electricity / dull looking [1 mark for each property up to a maximum of 2 marks].
 2.2 E.g. they can form more than one ion [1 mark], they make good catalysts [1 mark].
 2.3 Any two from: e.g. cobalt / copper / manganese / nickel / chromium / vanadium [1 mark for each correct answer]

Any two transition metals (apart from iron) will get you the marks.

Page 14 — Group 1 Elements

- 1.1 Y [1 mark]. As element Y has a higher melting point, it must be higher up the group than X [1 mark].

The higher up the group an element is, the lower its atomic number.

- 1.2 $2X_{(s)} + 2H_2O_{(l)} \rightarrow 2XOH_{(aq)} + H_{2(g)}$
 [1 mark for correct reactants and products and 1 mark for balanced equation. Half the ratio is acceptable]
 1.3 Anything between 8-14 [1 mark].
 2.1

	Boiling Point / °C	Radius of atom / pm
Rb	687.8	248
Cs	670.8	265
Fr	Accept lower than 670.8	Accept greater than 265

[1 mark for each correct answer]

- 2.2 Francium would be more reactive than caesium [1 mark]. As you go further down the group the outer electron is further away from the nucleus [1 mark], so the attraction between the nucleus and the electron decreases and the electron is more easily lost [1 mark].
 2.3 Formula: Fr_3P [1 mark]
 Equation: $12Fr + P_4 \rightarrow 4Fr_3P$ [1 mark for correct reactants and products, 1 mark for correctly balancing the equation]

Pages 15-16 — Group 7 Elements

Warm-up

Fluorine
 Chlorine
 Bromine
 Iodine

- 1.1 They are non-metals that exist as molecules of two atoms [1 mark].
 1.2 Chlorine is more reactive than bromine [1 mark]. This is because chlorine's outer shell is closer to the nucleus [1 mark] so it's easier for chlorine to gain an electron when it reacts [1 mark].

Because of the increasing distance between the nucleus and the outer shell, reactivity decreases down the group. Bromine is further down the group than chlorine, its outer shell is further away from the nucleus and therefore it's less reactive than chlorine.

- 1.3 P [1 mark]
 2.1 $2Fe + 3Br_2 \rightarrow 2FeBr_3$ [1 mark for Br_2 and 1 mark for balanced equation. Half the ratio is acceptable]
 2.2 -1 [1 mark]

All halide ions form ions with a -1 charge.

- 3.1 chlorine + potassium bromide \rightarrow potassium chloride + bromine [1 mark]
 3.2 The solution will turn orange [1 mark].
 3.3 displacement [1 mark]
 3.4 No [1 mark], as chlorine is less reactive than fluorine [1 mark].
 4.1 The halogens have seven electrons in their outer shell [1 mark]. As you go further down the group additional shells are added so the outer electron is further away from the nucleus [1 mark].
 4.2 Astatine will react more slowly than fluorine [1 mark] since reactivity decreases down the group [1 mark]. Both astatine and fluorine have 7 outer shell electrons so react in a similar way [1 mark]. So astatine will react with hydrogen to form hydrogen astatide/HAt [1 mark]. $H_2 + At_2 \rightarrow 2HAt$ [1 mark]


Page 17 — Group 0 Elements

- 1.1 Rn Boiling Point: Above $-108^\circ C$ [1 mark], Xe Density: Between 0.0037 and 0.0097 [1 mark], Ar Atomic Radius: Less than 109 pm [1 mark].
 1.2 Krypton is unreactive [1 mark]. It has a stable electron arrangement / full outer shell / 8 electrons in its outer shell [1 mark].
 1.3 Helium only has 2 electrons in its outer shell. The rest of the noble gases have 8 [1 mark].
 2.1 Noble gases are unreactive / they have stable electron arrangements / full outer shells / 8 electrons in their outer shell [1 mark].
 2.2 Iodine is much less reactive than fluorine [1 mark].
 2.3 Neon solidified at $-249^\circ C$ and xenon at $-112^\circ C$ [1 mark] Boiling points increase down the group [1 mark] and xenon is further down the group than neon so will have the higher boiling point [1 mark].

Topic 2 — Bonding, Structure and Properties of Matter

Page 18 — Formation of Ions

- 1.1 Metal atoms usually lose electrons to become positive ions [1 mark].



[2 marks if all four correct, otherwise 1 mark if two correct]


- 2.1 $2-$ [1 mark]
 2.2 2,8,8 [1 mark]. Sulfur gains two electrons [1 mark] to achieve a noble gas electronic structure/a full outer shell [1 mark].
 2.3 Argon/Ar [1 mark]

Pages 19-20 — Ionic Bonding

Warm-up

Dot and cross diagram	Ionic formula
	NaCl
	Na ₂ O
	MgCl ₂


1.1 calcium chloride [1 mark] and potassium oxide [1 mark]
Compounds that contain ionic bonding have to be made up of a metal and a non-metal. All the other options only contain non-metals, so can't be held together by ionic bonds.




[1 mark for arrow showing electron transfer from Li to F, 1 mark for correct electronic structure of fluoride ion, with seven crosses and one dot, 1 mark for correct charges on the ions.]

1.3 electrostatic attraction / electrostatic force [1 mark]

1.4 E.g. the particles in the compound are oppositely charged ions / have opposite charges / the bond is formed by electrons being transferred from one atom to another [1 mark].



If you showed the second electron shell of magnesium containing eight electrons as dots, you also get the mark.



2.2 E.g. the magnesium atom transfers two electrons to the oxygen atom [1 mark]. A magnesium ion with a 2+ charge forms [1 mark], and an oxide ion with a 2- charge forms [1 mark]. The oppositely charged ions are attracted to each other by electrostatic attraction [1 mark].

3.1 Element X: Group 7 [1 mark]

Reason: Any one of, e.g. it has formed an ion by gaining 1 electron / it forms 1- ions / the uncharged element would have seven electrons in its outer shell [1 mark].

Element Z: Group 2 [1 mark]

Reason: Any one of, e.g. it has formed an ion by losing 2 electrons / it forms 2+ ions / the uncharged element would have two electrons in its outer shell [1 mark].

3.2 How to grade your answer:

Level 0: There is no relevant information [No marks].

Level 1: The discussion is limited and doesn't mention both the uses and limitations of dot and cross diagrams [1 to 2 marks].

Level 2: There is some discussion of dot and cross diagrams, with at least one use and one limitation covered [3 to 4 marks].

Level 3: The discussion is comprehensive in evaluating both the uses and limitations of dot and cross diagrams [5 to 6 marks].

Here are some points your answer may include:

Dot and cross diagrams show:

Charge of the ions.

The arrangement of electrons in an atom or ion.

Which atoms the electrons in an ion originally come from.

Empirical formula (correct ratio of ions).

Dot and cross diagrams do not:

Show the structure of the compound.

Correctly represent the sizes of ions.

Pages 21-22 — Ionic Compounds

Warm-up

In an ionic compound, the particles are held together by **strong** forces of attraction. These forces act **in all directions** which results in the particles bonding together to form **giant lattices**.


1.1 conduct electricity in the solid state [1 mark]

1.2 giant ionic lattice [1 mark]

2.1 Sodium chloride contains positive sodium ions (Na⁺) [1 mark] and negative chloride ions (Cl⁻) [1 mark] that are arranged in a regular lattice/giant ionic lattice [1 mark]. The oppositely charged ions are held together by electrostatic forces acting in all directions [1 mark].

2.2 To melt sodium chloride, you have to overcome the very strong electrostatic forces/ionic bonds between the particles [1 mark], which requires lots of energy [1 mark].

3.1 E.g.



[1 mark for K⁺ ions, 1 mark for Br⁻ ions, 1 mark for correct structure, with alternating ions]

You'd also get the marks if you labelled all the white circles as Br⁻ and all the grey circles as K⁺.

3.2 Advantage: Any one of, e.g. the diagram shows the 3D arrangement of the ions / it suggests the structure is extended / it shows the regular (repeating) pattern of the ions [1 mark]. Disadvantage: Any one of, e.g. the diagram doesn't correctly represent the sizes of ions / it shows gaps between the ions [1 mark].

3.3 KBr [1 mark]

Remember that the overall charge of the ionic compound must be neutral. So you can work out the empirical formula by seeing that you only need one bromide ion to balance the charge on a potassium ion.

3.4 Boiling point: Potassium bromide has a giant structure with strong ionic bonds [1 mark]. In order to boil, these bonds need to be broken, which takes a lot of energy [1 mark].

Electrical conductivity of solid: The ions are in fixed positions in the lattice [1 mark] and so are not able to move and carry a charge through the solid [1 mark].

Electrical conductivity of solution: In solution, the ions are free to move [1 mark] and can carry a charge from place to place [1 mark].

Pages 23-24 — Covalent Bonding


1.1 They share a pair of electrons [1 mark].

1.2 Non-metals [1 mark]


1.3 BH₃ [1 mark]

Find the molecular formula by counting up how many atoms of each element there are in the diagram.


2



[1 mark]



[1 mark]



[1 mark]

Each line represents one covalent bond. Oxygen has a double bond, so you need to draw two lines between the oxygen atoms to show this.



3.1 E.g. it contains only non-metals [1 mark] and Figure 1 shows shared electrons [1 mark].

3.2 Any two from, e.g. they don't show how the atoms are arranged in space / they don't show the relative sizes of the atoms [2 marks — 1 mark for each correct answer].

- 3.3 One electron from hydrogen and one from carbon form a shared pair [1 mark] that are attracted to the nuclei of the carbon and hydrogen atoms [1 mark] by electrostatic attraction [1 mark].
- 4.1 Displayed formula: e.g. it shows how all the atoms in a molecule are connected in a simple way [1 mark], but it doesn't show the 3D structure of the molecule / it doesn't show which atom the electrons in the bond originally come from [1 mark].
Dot and cross diagram: e.g. it shows where the electrons in each covalent bond originally came from [1 mark] but it doesn't show the 3D structure of the molecule / they can become very complicated if the molecule is large [1 mark].
3D model: e.g. it shows how all the atoms are arranged in space in relation to each other / it shows the correct bond angles in the molecule [1 mark] but it quickly becomes complicated for large molecules / you can't tell which atom in the bonds the electrons originally came from [1 mark].
- 4.2 The displayed formula [1 mark] would be the best as it is easy to see how the atoms in a large molecule are connected without the diagram becoming too complicated [1 mark].

Pages 25-26 — Simple Molecular Substances


- 1.1 The bonds between the atoms are strong [1 mark], but the forces between the molecules are weak [1 mark].
- 1.2 The weak forces between the molecules / the intermolecular forces [1 mark].



- 2.3 E.g. N_2 has a triple covalent bond, whilst HCl has a single covalent bond [1 mark].
- 3.1 Simple molecular substances have weak forces between molecules [1 mark] so not much energy is needed to overcome them/they normally have low melting points [1 mark].
- 3.2 Iodine won't conduct electricity [1 mark] because the I_2 molecules aren't charged / the electrons aren't free to move so can't carry a charge [1 mark].
- 4.1 When methane boils, the forces between the molecules are overcome [1 mark] and it turns from a liquid into a gas [1 mark]. Methane is a smaller molecule than butane [1 mark] so the forces between the molecules are weaker [1 mark] and less energy is needed to overcome them [1 mark].
- 4.2 Carbon needs four more electrons to get a full outer shell, and does this by forming four covalent bonds [1 mark]. Hydrogen only needs one more electron to complete its outer shell, so can only form one covalent bond [1 mark].
Remember that the outer electron shell in hydrogen only needs two electrons to be filled, not eight like other electron shells.
- 4.3 Four [1 mark]. Silicon has four outer electrons so needs four more to get a full outer shell / silicon has the same number of outer shell electrons as carbon so will form the same number of bonds [1 mark].


Page 27 — Polymers and Giant Covalent Substances

Warm-up




- 1.1 Ammonia [1 mark]
Ammonia has a simple covalent structure — it forms small molecules.
- 1.2 The covalent bonds are very strong [1 mark], so a lot of energy is needed to break them [1 mark].
- 2.1 $(C_2H_4)_n$ [1 mark]
- 2.2 Solid [1 mark]. The molecule is very large and so the intermolecular forces are strong [1 mark] and need lots of energy to be broken [1 mark].
- 2.3 covalent bonds [1 mark]

Page 28 — Allotropes of Carbon

- 1.1 
- [2 marks if all correct, otherwise 1 mark if one correct]

- 1.2 A: graphene [1 mark]
B: buckminster fullerene [1 mark]
C: carbon nanotube / fullerene [1 mark]
- 1.3 Any one of, e.g. to strengthen materials / to deliver drugs into the body / as a catalyst / as a lubricant / in electronics [1 mark]
- 2.1 Graphite is made up of sheets of carbon atoms arranged in hexagons [1 mark], with weak forces between the sheets [1 mark]. Each carbon atom forms three covalent bonds [1 mark], and has one delocalised electron [1 mark].
- 2.2 Graphite has delocalised electrons [1 mark] which are free to move through the substance and carry an electric current [1 mark].

Page 29 — Metallic Bonding

- 1.1 E.g.  Metal ions in a regular pattern
Delocalised electrons
- [1 mark for regular arrangement of metal ions, 1 mark for delocalised electron, 1 mark for correct labels]
- 1.2 There is a strong electrostatic attraction [1 mark] between the delocalised electrons and the positive metal ions [1 mark].
- 1.3 High [1 mark] because the bonding is strong so requires lots of energy to break [1 mark].
- 1.4 Good [1 mark] because the electrons are free to move throughout the structure and carry an electrical charge [1 mark].
- 2.1 Metallic structures have layers of atoms [1 mark] that are able to slide over one another [1 mark].
- 2.2 Atoms of different elements are different sizes [1 mark]. Adding atoms of a different size to a pure metal distorts the layers [1 mark] making it harder for them to slide over one another [1 mark].

Page 30 — States of Matter

- 1.1 solid, liquid, gas [1 mark]
- 1.2 $NaCl_{(s)}$: solid [1 mark]
 $O_{2(g)}$: gas [1 mark]
 $Hg_{(l)}$: liquid [1 mark]
- 2.1 solid spheres [1 mark]
- 2.2 liquid [1 mark]
- 2.3 Any two from: melting / boiling / condensing / freezing [1 mark for each]
- 2.4 Any two from: e.g. the model says nothing about forces between particles / particles aren't really spheres / particles are mostly empty space, not solid [1 mark for each].


Page 31 — Changing State

- 1.1 melting [1 mark]
 1.2 boiling point [1 mark]
 1.3 The bonds are strong [1 mark].
 2.1 sodium chloride [1 mark]
 At 900 °C, water would be a gas and copper would be a solid.
 2.2 Sodium chloride [1 mark] and water [1 mark].
 At 1500 °C, copper would be a liquid.
 2.3 Boiling sodium chloride [1 mark].
 2.4 No [1 mark]. When copper boils, the metallic bonds are broken [1 mark], but when water boils only the intermolecular forces are broken [1 mark], so you can't tell anything about the strength of the covalent bonds [1 mark].

Pages 32-33 — Nanoparticles

Warm-up

- 1 True
 2 False
 3 True
 4 True
 5 False



[2 marks if all three correct, otherwise 1 mark if one correct]

- 2.1 $10 \text{ nm} \times 10 \text{ nm} \times 10 \text{ nm} = 1000 \text{ nm}^3$ [1 mark]
 2.2 Area of one side = $10 \text{ nm} \times 10 \text{ nm} = 100 \text{ nm}^2$
 A cube has six sides, so total surface area = $6 \times 100 \text{ nm}^2 = 600 \text{ nm}^2$ [2 marks for correct answer, otherwise 1 mark for the area of one side]
 2.3 Surface area to volume ratio = $600 \text{ nm}^2 \div 1000 \text{ nm}^3 = 0.6 \text{ nm}^{-1}$ [1 mark]
 2.4 Volume = $1 \text{ nm} \times 1 \text{ nm} \times 1 \text{ nm} = 1 \text{ nm}^3$
 Surface area of one side = $1 \text{ nm} \times 1 \text{ nm} = 1 \text{ nm}^2$
 Total surface area of cube = $6 \times 1 \text{ nm}^2 = 6 \text{ nm}^2$
 Surface area to volume ratio = $6 \text{ nm}^2 \div 1 \text{ nm}^3 = 6 \text{ nm}^{-1}$
 [4 marks for correct answer, otherwise 1 mark for correct volume, 1 mark for correct surface area of one side, 1 mark for correct total surface area]
 E.g. decreasing the side length by a factor of 10 increases the surface area to volume ratio by a factor of 10 [1 mark].
 3.1 Surface area to volume ratio = $0.12 \div 10 = 0.012 \text{ nm}^{-1}$ [1 mark]
 The side length has increased by a factor of ten, so the surface area to volume ratio will decrease by a factor of 10.
 3.2 Material Y [1 mark], because the particles have a lower surface area to volume ratio [1 mark].


Page 34 — Uses of Nanoparticles

- 1.1 E.g. suncreams containing nanoparticles give better skin coverage [1 mark] and are better at absorbing UV rays [1 mark].
 1.2 E.g. the nanoparticles in suncreams may affect people's health [1 mark], and when they're washed away they could damage the environment [1 mark].
 2.1 Material: carbon nanotubes [1 mark]
 Reason: e.g. the carbon nanotubes trap the drug molecules inside, and release them when they're at the right place in the body [1 mark].
 2.2 Material: silver [1 mark]
 Reason: e.g. silver nanoparticles are antibacterial, so will help to kill any bacteria that are in the water [1 mark].

- 2.3 Material: carbon nanotubes [1 mark]
 Reason: e.g. carbon nanotubes are strong, so will strengthen the sports equipment, but they are also light, so won't add much weight to the equipment [1 mark].

Topic 3 — Quantitative Chemistry

Page 35 — Relative Formula Mass



- 2.1 $M_r(\text{MgO}) = 24 + 16 = 40$ [1 mark]
 percentage by mass of magnesium = $\frac{A_r(\text{Mg})}{M_r(\text{MgO})} \times 100 = \frac{24}{40} \times 100 = 60\%$ [1 mark]
 2.2 Mass of magnesium ions = $200 \times \frac{15}{100} = 30 \text{ g}$ [1 mark]
 2.3 Mass of magnesium oxide containing 30 g of magnesium ions = $30 \div \frac{60}{100} = 50 \text{ g}$ [1 mark]

If you used the percentage mass of magnesium ions as 40% and the mass of magnesium ions in the mixture as 20 g, your answer will also be 50 g.

Page 36 — The Mole

Warm-up:

- 6.02×10^{23}
 1.1 M_r of carbon dioxide = $12 + (16 \times 2) = 44$ [1 mark]
 1.2 Moles of carbon dioxide = $110 \div 44 = 2.5 \text{ mol}$ [1 mark]
 1.3 1 mole of carbon dioxide would weigh more [1 mark]. It has a higher relative formula mass [1 mark].
 2.1 2 mol sulfur = $2 \times 32 \text{ g} = 64 \text{ g}$ [1 mark]
 2.2 M_r of iron sulfide = $56 + 32 = 88$
 Moles of iron sulfide = $44 \div 88 = 0.50 \text{ mol}$ [2 marks for correct answer, otherwise 1 mark for correct working]
 2.3 The number of atoms in 3 moles of sulfur is greater than the number of molecules in 2 moles of iron sulfide [1 mark]. There's the same number of atoms in 1 mole of sulfur as there are molecules in 1 mole of iron sulfide so in 3 moles of sulfur there will be more atoms than there are molecules in 2 moles of iron sulfide [1 mark].

Pages 37-38 — Conservation of Mass

- 1.1 $2\text{Mg} + \text{O}_2 \rightarrow 2\text{MgO}$ [1 mark]
 1.2 Mass of oxygen = 20 g of MgO – 12 g of Mg = 8 g [2 marks for correct answer, otherwise 1 mark for correct working]
 2.1 The mass of reactants equals the mass of products in a chemical reaction [1 mark]. Atoms are not made or destroyed during a chemical reaction [1 mark]. So, there must be the same number of each type of atom in the products as in the reactants [1 mark].
 2.2 The mass of the powder would increase [1 mark]. Oxygen gas was not included as part of the original measurement [1 mark]. Particles of oxygen are added to the zinc to form zinc oxide powder [1 mark].
 3.1 The measurement is correct [1 mark]. Carbon dioxide (a gas) is produced and released into the atmosphere [1 mark]. So, the student only measured the mass of the solid product, not both reactants [1 mark].
 3.2 M_r of sodium oxide = $106 - 44 = 62$ [1 mark]
 3.3 Moles of $\text{Na}_2\text{CO}_3 = 53 \div 106 = 0.50$
 For every mole of Na_2CO_3 that reacts, 1 mole of CO_2 is produced. Only 0.50 moles of Na_2CO_3 react so 0.50 moles of CO_2 are produced.
 Mass of carbon dioxide = $0.50 \times 44 = 22 \text{ g}$ [3 marks for correct answer, otherwise 1 mark for 0.50 moles of Na_2CO_3 and 1 mark for a 1:1 molar ratio]

To work out a molar ratio, you need to use the balanced symbol equation


for the reaction. The numbers in front of the chemical formulas show the number of moles of a substance that react or are produced in the reaction. In this question, for every 1 mole Na_2CO_3 heated, 1 mole of carbon dioxide is produced — a 1:1 molar ratio.

3.4 Mass of sodium oxide = $53 \text{ g} - 22 \text{ g} = 31 \text{ g}$ [1 mark]

Pages 39-40 — The Mole and Equations

Warm-up

3



2.1 Moles of sodium = $9.2 \div 23 = 0.4 \text{ mol}$ [1 mark]

2.2 M_r of water = $(1 \times 2) + 16 = 18$

Moles of water = $7.2 \text{ g} \div 18 = 0.4 \text{ mol}$ [2 marks for correct answer, otherwise 1 mark for correct working]


2.3 Divide the number of moles of each substance by the lowest of these number of moles (0.2 mol) to give the molar ratios.

$\text{Na} = 0.4 \div 0.2 = 2 \text{ mol}$

$\text{H}_2\text{O} = 0.4 \div 0.2 = 2 \text{ mol}$

$\text{NaOH} = 0.4 \div 0.2 = 2 \text{ mol}$

$\text{H}_2 = 0.2 \div 0.2 = 1 \text{ mol}$



3.1 Moles of methane = $8 \text{ g} \div 16 = 0.5 \text{ mol}$

Moles of oxygen = $32 \text{ g} \div 32 = 1 \text{ mol}$

Moles of carbon dioxide = $22 \text{ g} \div 44 = 0.5 \text{ mol}$

Moles of water = $18 \text{ g} \div 18 = 1 \text{ mol}$ [1 mark]


Divide by the lowest of these numbers which is 0.5:

Methane = $0.5 \div 0.5 = 1 \text{ mol}$

Oxygen = $1 \div 0.5 = 2 \text{ mol}$

Carbon dioxide = $0.5 \div 0.5 = 1 \text{ mol}$

Water = $1 \div 0.5 = 2 \text{ mol}$ [1 mark]



3.2 Moles of oxygen = $48 \text{ g} \div 32 = 1.5 \text{ mol}$

Molar ratio of oxygen : carbon dioxide = 2:1

Moles of carbon dioxide = $1.5 \text{ mol} \div 2 = 0.75 \text{ mol}$ [3 marks for correct answer, otherwise 1 mark for 1.5 mol of oxygen and 1 mark for molar ratio of 2:1]

3.3 Molar ratio of CH_4 : $\text{H}_2\text{O} = 1:2$

4 mol of methane will produce 8 mol of water [1 mark].

3.4 Mass of water = $18 \times 8 = 144 \text{ g}$ [1 mark]

If you got the equation wrong in 3.1 but used all the right working in parts 3.2, 3.3 and 3.4, you still get the marks, even if you got a different answer to the one here.

Page 41 — Limiting Reactants

1.1 To make sure that all the hydrochloric acid was used up in the reaction [1 mark].

1.2 The limiting reactant is completely used up during a reaction [1 mark] and so its quantity limits the amount of product that can be formed [1 mark].

2.1 Molar ratio of copper oxide : copper sulfate = 1:1

Therefore, 0.50 mol of copper sulfate is produced.

M_r of copper sulfate = $63.5 + 32 + (16 \times 4) = 159.5$

Mass of copper sulfate = $0.50 \times 159.5 = 80 \text{ g}$ [3 marks for correct answer, otherwise 1 mark for 0.50 moles of copper sulfate and 1 mark for M_r of 159.5]

2.2 The amount of product formed is directly proportional to the amount of limiting reactant [1 mark]. So doubling the quantity of the sulfuric acid will double the yield of the copper sulfate [1 mark].

2.3 If only 0.4 mol of copper oxide is present, there will not be enough molecules to react with all the sulfuric acid [1 mark]. The copper oxide will be the limiting reactant [1 mark] and only 0.4 mol of product will be formed [1 mark].

Pages 42-43 — Gases and Solutions

1.1 Conc. of calcium chloride = $28 \text{ g} \div 0.4 \text{ dm}^3 = 70 \text{ g/dm}^3$
[1 mark for correct answer and 1 mark for correct units]

1.2 The concentration of a solution is the amount of a substance in a given volume of a solution [1 mark].

2.1 Mass of $\text{CO}_2 = (36 + 24) \times 44 = 66 \text{ g}$

[2 marks for correct answer, otherwise 1 mark for using the correct equation to calculate mass]

2.2 1 mole of any gas at room temperature and pressure has a volume of 24 dm^3 [1 mark]. So, 1 mole of CO_2 and 1 mole of O_2 will have the same volume [1 mark].

3.1 M_r of $\text{O}_2 = 2 \times 16 = 32$

Volume of oxygen = $(16 \text{ g} \div 32) \times 24 = 12 \text{ dm}^3$

[3 marks for correct answer, otherwise 1 mark for M_r of oxygen, 1 mark for using the correct equation to calculate volume]

3.2 Molar ratio of oxygen : carbon dioxide = 2:1

Volume of carbon dioxide = $12 \div 2 = 6 \text{ dm}^3$ [2 marks for correct answer, otherwise 1 mark for 2:1 molar ratio]

4.1 Volume of oxygen = $48 \div 2 = 24 \text{ dm}^3$ [1 mark]

As the molar ratio of carbon monoxide to oxygen is 2:1, there must be half the volume of oxygen as there is carbon monoxide.

4.2 Volume of carbon monoxide: $(28 \div 28) \times 24 = 24 \text{ dm}^3$

Carbon monoxide to oxygen molar ratio = 2:1 so volume of oxygen = $24 \div 2 = 12 \text{ dm}^3$ [4 marks for correct answer, otherwise 1 mark for using the formula

volume = mass $\div M_r \times 24$, 1 mark for correct volume of carbon monoxide and 1 mark for 2:1 molar ratio]

4.3 24 dm^3 [1 mark]

The molar ratio of carbon monoxide to carbon dioxide is 2:2, therefore there is the same volume of carbon dioxide as there is carbon monoxide.

5.1 Moles of $\text{Na}_2\text{CO}_3 = 0.50 \text{ mol/dm}^3 \times 0.50 \text{ dm}^3 = 0.25 \text{ mol}$ [1 mark]

5.2 M_r of $\text{Na}_2\text{CO}_3 = (23 \times 2) + 12 + (16 \times 3) = 106$

Mass of 0.25 mol of $\text{Na}_2\text{CO}_3 = 0.25 \times 106 = 26.5 \text{ g}$ [2 marks for correct answer, otherwise 1 mark for correct working]

Pages 44-45 — Concentration Calculations

1.1 M_r of HCl = $1 + 35.5 = 36.5$

Concentration of HCl = $18.25 \div 36.5 = 0.500 \text{ mol/dm}^3$

[3 marks for correct answer, otherwise 1 mark for M_r of HCl, 1 mark for correct equation to convert the units of concentration]

1.2 Volume of HCl = $25.0 \text{ cm}^3 \div 1000 = 0.0250 \text{ dm}^3$

Moles of HCl = $0.500 \text{ mol/dm}^3 \times 0.0250 \text{ dm}^3 = 0.0125 \text{ mol}$

[3 marks for correct answer, otherwise 1 mark for concentration of HCl in dm^3 and 1 mark for correct equation to calculate moles]

1.3 Moles of NaOH = 0.0125 mol [1 mark]

The molar ratio of hydrochloric acid to sodium hydroxide is 1:1, so there must be the same number of moles of sodium hydroxide as there are hydrochloric acid.

1.4 Volume of NaOH = $50.0 \text{ cm}^3 \div 1000 = 0.0500 \text{ dm}^3$

Concentration of NaOH = $0.0125 \text{ mol} \div 0.0500 \text{ dm}^3$


$= 0.250 \text{ mol/dm}^3$

[3 marks for correct answer, otherwise 1 mark for volume of NaOH in dm^3 and 1 mark for correct equation to calculate concentration]

1.5 M_r of NaOH = $23 + 16 + 1 = 40$

Concentration of NaOH = $0.250 \text{ mol} \times 40 = 10 \text{ g/dm}^3$

[3 marks for correct answer, otherwise 1 mark for M_r of NaOH and 1 mark for correct equation to convert the units of concentration]



[1 mark for correct symbols, 1 mark for correct balancing]

2.2 Mean volume of HCl = $(12.50 + 12.55 + 12.45) \div 3$

$= 12.50 \text{ cm}^3$

[2 marks for correct answer, otherwise 1 mark for correct equation to calculate the mean]

2.3 Volume of HCl = $12.50 \text{ cm}^3 \div 1000 = 0.01250 \text{ dm}^3$

Volume of $\text{Na}_2\text{CO}_3 = 25.0 \text{ cm}^3 \div 1000 = 0.0250 \text{ dm}^3$

Moles of HCl = $0.0125 \text{ dm}^3 \times 1.00 \text{ mol/dm}^3 = 0.0125 \text{ mol}$

Molar ratio of HCl : $\text{Na}_2\text{CO}_3 = 2:1$

Moles of $\text{Na}_2\text{CO}_3 = 0.01250 \text{ mol} \div 2 = 0.006250 \text{ mol}$
 Concentration of $\text{Na}_2\text{CO}_3 = 0.006250 \text{ mol} \div 0.0250 \text{ dm}^3 = 0.250 \text{ mol/dm}^3$

[6 marks for correct answer, otherwise 1 mark for volumes of HCl and Na_2CO_3 in dm^3 , 1 mark for correct equation to calculate moles, 1 mark for moles of HCl, 1 mark for moles of Na_2CO_3 and 1 mark for correct equation to calculate concentration]

If answer to question 2.2 is incorrect, but your working is correct here, you still get all the marks, even if you got a different answer.

Page 46 — Atom Economy

- 1.1 Any two from: e.g. less waste / more sustainable / more profitable [2 marks — 1 mark for each correct answer].
- 1.2 M_r of ethanol = $(12 \times 2) + (1 \times 6) + 16 = 46$ [1 mark]
- 1.3 M_r of ethene = $(12 \times 2) + (1 \times 4) = 28$ [1 mark]
- 1.4 Atom economy = $(28 \div 46) \times 100 = 61\%$
 [2 marks for correct answer, otherwise 1 mark for correct equation for atom economy]
- 2 Atom economy of reaction using magnesium:
 M_r of reactants = $A_r(\text{Mg}) + (2 \times M_r(\text{HCl}))$
 $= 24 + (2 \times 36.5) = 97$
 Atom economy = $(2 \div 97) \times 100 = 2\%$ [2 marks for correct answer, otherwise 1 mark for M_r of reactants]
 Atom economy of reaction using zinc:
 M_r of reactants = $A_r(\text{Zn}) + (2 \times M_r(\text{HCl}))$
 $= 65 + (2 \times 36.5) = 138$
 Atom economy = $(2 \div 138) \times 100 = 1\%$ [2 marks for correct answer, otherwise 1 mark for M_r of reactants]
 More economical reaction: the magnesium reaction [1 mark]

Page 47 — Percentage Yield

- 1.1 Percentage yield = $(1.8 \text{ g} \div 2.4 \text{ g}) \times 100 = 75\%$
 [2 marks for correct answer, otherwise 1 mark for correct method]
- 1.2 Any one from: e.g. some of the magnesium may not yet have reacted / some product may have been left behind in the crucible [1 mark].
- 2.1 Mass of N_2 in g = $14 \times 1000 = 14\,000 \text{ g}$
 Number of moles of $\text{N}_2 = 14\,000 \div (2 \times 14) = 500 \text{ mol}$
 500 mol of N_2 react to produce $2 \times 500 = 1000 \text{ mol}$ of NH_3 .
 M_r of $\text{NH}_3 = 14 + (3 \times 1) = 17$
 Theoretical yield = moles $\times M_r = 1000 \times 17 = 17\,000 \text{ g} = 17 \text{ kg}$
 [4 marks for correct answer, otherwise 1 mark for correct number of moles of N_2 , 1 mark for correct number of moles of NH_3 and 1 mark for M_r of NH_3]
- 2.2 Percentage yield = $(4.5 \text{ kg} \div 17 \text{ kg}) \times 100 = 26\%$
 [2 marks for correct answer, otherwise 1 mark for correct method]
- 2.3 Any two from: e.g. the reaction is reversible so may not have gone to completion / products may have been lost during the reaction / there may have been side reactions [1 mark for each correct answer].
- 2.4 Any two from: e.g. to reduce waste / increase sustainability / to reduce cost [2 marks — 1 mark for each correct answer].

Topic 4 — Chemical Changes

Page 48 — Acids and Bases

Warm-up


Universal indicator will turn **red** in strongly acidic solutions and **purple** in strongly alkaline solutions. In a **neutral** solution, Universal indicator will be green. A pH probe attached to a pH meter is **more** accurate than Universal indicator as it displays a numerical value for pH.

- 1.1 beer [1 mark]
 1.2 blue / blue-green [1 mark]
 1.3 H^+ [1 mark]
 1.4 0 [1 mark] – 14 [1 mark]

- 2.1 acid + alkali \rightarrow salt + water [1 mark]
 2.2 $\text{H}^+_{(\text{aq})} + \text{OH}^-_{(\text{aq})} \rightarrow \text{H}_2\text{O}_{(\text{l})}$ [1 mark]
 You still get the marks if you didn't include state symbols.

Page 49 — Titrations

Warm-up



- 1.1 Universal indicator is the most suitable indicator for use in titrations [1 mark].
- 1.2 A burette allows an acid/alkali to be added to a solution drop-by-drop [1 mark] which helps determine the end-point more accurately [1 mark].
- 1.3 The titration should be repeated several times to achieve several consistent readings [1 mark]. The mean reading should be used to calculate the concentration [1 mark].

Page 50 — Strong Acids and Weak Acids

- 1.1 A strong acid completely ionises/dissociates in solution [1 mark]. A weak acid only partly ionises in solution [1 mark].
- 1.2 Nitric acid would have a lower pH than ethanoic acid [1 mark] because it is a stronger acid/more dissociated/ionised [1 mark], so the concentration of H^+ would be greater [1 mark].

You would also get the marks for using the reverse argument — ethanoic acid would have a higher pH because it is a weaker acid so the concentration of H^+ ions is lower.

- 1.3 3 [1 mark]
 As the concentration of H^+ ions in solution decreases by a factor of 10, the pH rises by 1.
- 1.4 Adding water to the beaker [1 mark].
 Adding ethanoic acid to the beaker at the same concentration as the citric acid [1 mark].
 Changing the citric acid to carbonic acid of the same concentration [1 mark].

Pages 51-52 — Reactions of Acids

- 1.1 Neutralisation [1 mark]
 1.2 Fizzing — Carbon dioxide is produced [1 mark]
 2.1 sulfuric acid + lithium hydroxide \rightarrow lithium sulfate + water [1 mark]
 2.2 $\text{H}_2\text{SO}_4 + 2\text{LiOH} \rightarrow \text{Li}_2\text{SO}_4 + 2\text{H}_2\text{O}$ [1 mark for correct formula of Li_2SO_4 , 1 mark for correct balancing]
 2.3 Both reactions produce lithium sulfate and water [1 mark]. The reaction between sulfuric acid and lithium carbonate also produces carbon dioxide [1 mark].
- 3.1 Add zinc oxide to hydrochloric acid until the reaction stops / the excess metal oxide sinks to the bottom [1 mark]. Filter the excess solid from the solution using a filter funnel [1 mark]. Heat the zinc chloride solution to evaporate some of the water and then leave to cool [1 mark]. Filter and dry the crystals that form [1 mark].
- 3.2 E.g. zinc carbonate [1 mark].
 Any other insoluble zinc base or zinc metal also gets a mark.
- 4 How to grade your answer:
 Level 0: Nothing written worthy of credit [No marks].
 Level 1: Some suitable tests are named but it is not clear how the results would enable the solutions to be identified. The chemistry of the tests is not clearly described [1 to 2 marks].
 Level 2: Tests that enable at least one solution to be identified are clearly described, or tests that would enable all solutions to be identified are named but not clearly described [3 to 4 marks].

Level 3: At least two tests are described together with the expected outcomes. It is clear how these tests would be used to distinguish between all three solutions. The chemistry of the tests is correctly described [5 to 6 marks].

Here are some points your answer may include:

Test the pH of each solution.

The neutral solution/the solution that turns Universal indicator green is the salt.

Add a couple of drops of Universal indicator to the solutions followed by some dilute acid.

The solution containing sodium carbonate will fizz as it reacts with the acid to release carbon dioxide gas as shown by the equation: acid + sodium carbonate → sodium salt + water + carbon dioxide

The solution containing sodium hydroxide will react with acid changing the Universal indicator solution from blue/purple to green, but there won't be any fizzing as no gas is released as shown by the reaction:

acid + sodium hydroxide → sodium salt + water

The solution containing the sodium salt won't react with acid.

Pages 53-54 — The Reactivity Series

- 1.1 magnesium + hydrochloric acid → magnesium chloride + hydrogen [1 mark]
- 1.2 Positive magnesium ions [1 mark]
- 1.3 It forms positive ions less easily / it's lower down in the reactivity series [1 mark].
- 1.4 Any one of: e.g. potassium / sodium / lithium / calcium [1 mark].
- 2.1 metal + water → metal hydroxide + hydrogen [1 mark]
- 2.2 $\text{Ca}_{(s)} + 2\text{H}_2\text{O}_{(l)} \rightarrow \text{Ca}(\text{OH})_{2(aq)} + \text{H}_{2(g)}$ [1 mark for each correct product]
- 2.3 Any one from: e.g. lithium / sodium / potassium [1 mark]
As it is higher in the reactivity series than calcium / loses electrons more easily than calcium / forms positive ions more easily [1 mark].
- 2.4 potassium, sodium, zinc [1 mark]
- 3.1 When a metal reacts with an acid, the metal forms positive ions [1 mark]. The results show that lithium reacts more vigorously with acid than magnesium does [1 mark], so lithium forms positive ions more easily [1 mark].
- 3.2 A very vigorous fizzing/more vigorous than lithium [1 mark], sodium disappears [1 mark].
- 3.3 lithium, calcium, copper [1 mark]
- 3.4 It is not possible to tell the difference between magnesium and zinc from these results since both have same reaction with dilute acid [1 mark]. E.g. to find which is more reactive, you could find the effect of adding zinc to water [1 mark].

Page 55 — Separating Metals from Metal Oxides

- 1.1 E.g. gold [1 mark]
- 1.2 Many metals can react with other elements/oxygen to form compounds/oxides [1 mark].
- 1.3 Reduction is the loss of oxygen [1 mark].
- 1.4 Magnesium is more reactive than carbon [1 mark].
- 2.1 $2\text{Fe}_2\text{O}_3 + 3\text{C} \rightarrow 4\text{Fe} + 3\text{CO}_2$
[1 mark for correct equation, 1 mark for correct balancing]
- 2.2 Carbon has been oxidised [1 mark] as it has gained oxygen during this reaction [1 mark].
- 2.3 E.g. extracting magnesium would have high energy costs to provide the high temperature and reduced pressure needed [1 mark], but iron extraction doesn't need to be continuously heated [1 mark].

Page 56 — Redox Reactions

- 1.1 Reduction is the gain of electrons [1 mark].
- 1.2 zinc chloride + sodium → zinc + sodium chloride [1 mark]
- 1.3 Hydrogen gains electrons [1 mark].

1.4 Chlorine is neither oxidised nor reduced [1 mark].


2.1 $\text{Mg}_{(s)} + \text{Fe}^{2+}_{(aq)} \rightarrow \text{Mg}^{2+}_{(aq)} + \text{Fe}_{(s)}$ [1 mark]

You still get the marks if you didn't include state symbols.


2.2 No reaction would occur [1 mark]. Copper is less reactive than iron so doesn't displace it [1 mark].

Pages 57-59 — Electrolysis

Warm-up



- 1.1 A liquid or solution that can conduct electricity [1 mark].
- 1.2 lead bromide → lead + bromine [1 mark]
- 1.3 Lead ions have a positive charge [1 mark]. This means they are attracted to the negative cathode [1 mark].
- 1.4 Br⁻ [1 mark]
- 1.5 oxidation [1 mark]
- 1.6 So the ions can move to the electrodes [1 mark].
- 2.1 molten aluminium [1 mark]
- 2.2 To lower the melting point of the electrolyte [1 mark].
- 2.3 Carbon in the electrodes reacts with oxygen to form carbon dioxide [1 mark], so they degrade over time [1 mark].
- 3.1 Iron ions, chloride ions, hydrogen ions and hydroxide ions [1 mark for iron ions and chloride ions, 1 mark for hydrogen ions and hydroxide ions].
- 3.2 At the cathode: hydrogen is discharged.
- 3.3 At the anode: chlorine is discharged [1 mark].
- 3.4 oxygen [1 mark]
- 3.4 Iron can be extracted via reduction with carbon [1 mark], which is less expensive than electrolysis [1 mark].
- 4.1 E.g.



[1 mark for power supply, 1 mark for electrodes in solution, 1 mark for boiling tubes over the electrodes, 1 mark for labels]

4.2

Solution	Product at cathode	Product at anode
CuCl_2	Cu	Cl_2
KBr	H_2	Br_2
H_2SO_4	H_2	O_2 and H_2O

[1 mark for each correct answer]

- 4.3 Potassium is more reactive than hydrogen [1 mark] so hydrogen is discharged [1 mark]. There are no halide ions [1 mark] so oxygen and water are discharged [1 mark].
- 4.4 Cathode: $2\text{H}^+ + 2\text{e}^- \rightarrow \text{H}_2$ [1 mark]
Anode: $4\text{OH}^- \rightarrow \text{O}_2 + 2\text{H}_2\text{O} + 4\text{e}^-$
/ $4\text{OH}^- - 4\text{e}^- \rightarrow \text{O}_2 + 2\text{H}_2\text{O}$
[1 mark]


Topic 5 — Energy Changes

Pages 60-61 — Exothermic and Endothermic Reactions

1 In an endothermic reaction, energy is transferred from the surroundings so the temperature of the surroundings goes down [1 mark].

2.1 endothermic [1 mark]

2.2



[1 mark for correct curve, 1 mark for energy change]

The curve has to go above the energy of the products and then fall back down. If you didn't do this, you don't get the mark.

2.3 From the surroundings [1 mark].

2.4 It stays the same [1 mark].

2.5 E.g. a sports injury pack [1 mark].

3.1 The activation energy is the minimum amount of energy that reactants must have when they collide with each other in order to react [1 mark]. It's shown by the difference between the energy of the reactants and the maximum energy reached by the curve on the reaction profile [1 mark].

3.2 Reaction A is the most suitable reaction [1 mark].

Reaction C is endothermic, so would not give out heat, and couldn't be used to warm your hands [1 mark].

Reaction A has a lower activation energy than Reaction B / gives out more energy than Reaction B [1 mark].

4.1 Any three from: e.g. thermometer / polystyrene cup (and lid) / mass balance / measuring cylinder / beaker filled with cotton wool / stopwatch [1 mark for each].

4.2 How to grade your answer:

Level 0: There is no relevant information [No marks].

Level 1: The method is vague, and misses out important details about how the investigation could be carried out [1 to 2 marks].

Level 2: The method is clear, but misses out a few key details about how the investigation would be carried out or how the variables could be controlled [3 to 4 marks].

Level 3: There is a clear and detailed method that includes ways to reduce energy transfer to the surroundings, and specifies variables that should be controlled throughout the investigation [5 to 6 marks].

Here are some points your answer may include:

Measure out an exact volume of the acid solution into the polystyrene cup.

Record the initial temperature of the acid solution.

Add one metal powder and stir the mixture.

Place a lid on the polystyrene cup to reduce the amount of energy transferred to the surroundings.

Take the temperature of the mixture every 30 seconds and record the highest temperature.

Repeat the experiment for each different metal.

Use the same volume and concentration of acid each time you repeat the experiment.

Make sure the acid starts at the same temperature each time you repeat the experiment.

Use the same number of moles and the same surface area of metal each time you repeat the experiment.

Page 62 — Bond Energies

1.1 Energy to break the bonds = $(4 \times \text{C-H}) + \text{Cl-Cl}$
 $= (4 \times 413) + 243 = 1652 + 243 = 1895 \text{ kJ/mol}$
 Energy produced when bonds form = $(3 \times \text{C-H}) + \text{C-Cl} + \text{H-Cl}$
 $= (3 \times 413) + 346 + 432 = 1239 + 346 + 432$
 $= 2017 \text{ kJ/mol}$

Energy change of reaction = Energy to break bonds – Energy produced when bonds form

$= 1895 - 2017 = -122 \text{ kJ/mol}$ [3 marks for correct answer, otherwise 1 mark for 1895 kJ/mol, 1 mark for 2017 kJ/mol, 1 mark for subtracting energy produced when bonds form from energy needed to break bonds]

Three of the C-H bonds are unchanged in this reaction. So you could also calculate this by working out just the energy needed to break the C-H and the Cl-Cl bond, and subtracting the energy that's released when the new C-Cl and H-Cl bonds form.

1.2 The reaction is exothermic [1 mark] because the energy released when the bonds of the products form is greater than the energy needed to break the bonds of the reactants [1 mark].

2 Total energy needed to break the bonds in the reactants
 $= \text{H-H} + \text{F-F} = 436 + 158 = 594 \text{ kJ/mol}$
 Energy change of reaction = Energy needed to break bonds – Energy released when bonds form

So, energy released when bonds form = Energy needed to break bonds – Energy change of reaction

$= 594 - (-542) = 1136 \text{ kJ/mol}$

Energy released when bonds form = $2 \times \text{H-F}$ bond energy

So, H-F bond energy = $1136 \div 2 = 568 \text{ kJ/mol}$

[3 marks for correct answer, otherwise 1 mark for finding the energy needed to break the bonds, 1 mark for finding the energy released by forming bonds.]

Page 63-64 — Cells, Batteries and Fuel Cells


Warm-up

1 False

2 False

3 True

1



[2 marks for three correct lines, 1 mark for one correct line]

2.1 Cell B [1 mark] as if the two metals are the same then no voltage will be produced / the two metals must be different in order for a voltage to be produced [1 mark].

2.2 The set-up is a battery [1 mark]. The voltage will increase to be twice the voltage of Cell B by itself [1 mark].

2.3 The reactants/chemicals in the cells get used up [1 mark], so a voltage is no longer produced [1 mark].

2.4 It can be reversed by connecting the cell to an external electric current [1 mark].

3.1 Reaction equation: $\text{H}_2 \rightarrow 2\text{H}^+ + 2\text{e}^-$ [1 mark]

Type of reaction: oxidation [1 mark]

3.2 water [1 mark]

3.3 Any one of: e.g. hydrogen fuel cells are less polluting to dispose of than rechargeable batteries / there's a limit to how many times a rechargeable battery can be recharged, but this isn't a problem for hydrogen fuel cells / fuel cells store more energy than rechargeable batteries [1 mark].

4.1 C, A, D, B [2 marks, or 1 mark if 2 correct]

The greater the voltage of the cell, the more reactive the metal [1 mark]. Metal C produces the greatest voltage, so is the most reactive, followed by A, then D, and finally B which produces the lowest voltage [1 mark].


4.2 E.g. the electrolyte [1 mark].

Topic 6 — The Rate and Extent of Chemical Change

Pages 65-67 — Rates of Reaction


1.1 Using a larger volume of the solution, but keeping the concentration the same [1 mark].

- 1.2 activation energy [1 mark]
- 1.3 A catalyst decreases the activation energy [1 mark].
- 2 Produced most product: C [1 mark]
- Finished first: B [1 mark]
- Started at the slowest rate: A [1 mark]



[1 mark for curve with steeper gradient at the start of the reaction, 1 mark for curve reaching the final volume earlier, 1 mark for final volume being the same as for the other curve]

- 3.2 The frequency of the collisions [1 mark] and the energy of the colliding particles [1 mark].
- 3.3 There are more particles in a given volume/the particles are closer together [1 mark], so the collisions between particles are more frequent [1 mark].
- 3.4 The rate would increase [1 mark].
- 3.5 Smaller pieces have a higher surface area to volume ratio [1 mark]. So for the same volume of solid, the particles around it will have more area to work on and collisions will be more frequent [1 mark].
- 3.6 E.g. changing the temperature / adding a catalyst [1 mark].
- 4.1 E.g. increasing the volume of the reaction vessel would decrease the pressure of the reacting gases [1 mark]. So the particles would be more spread out and would collide less frequently [1 mark], so the reaction rate would decrease [1 mark]. Increasing the temperature would cause the particles to move faster, so the frequency of collisions would increase [1 mark] and the reaction rate would increase [1 mark].
- 4.2 It's a catalyst [1 mark].
- 4.3 The reaction equation won't change [1 mark]. Cerium oxide isn't used up in the reaction, so doesn't appear in the reaction equation [1 mark].



[1 mark for correct relative energies of products and reactants, 1 mark for start and end energies being the same for reactions with and without cerium oxide, 1 mark for reaction with cerium oxide rising to a lower energy than reaction without cerium oxide]


Pages 68-70 — Measuring Rates of Reaction

Warm-up

The rate of a reaction can be measured by dividing the amount of **reactants** used up or the amount of **products** formed by the **time**. To find the rate at a particular time from a graph with a curved line of best fit, you have to find the **gradient** of the **tangent** at that time.

- 1 mass [1 mark], volume of gas [1 mark]
- 2.1 time taken for the solution to go cloudy [1 mark]

- 2.2 temperature [1 mark]
- 2.3 Any one from: e.g. the concentration of the reactants / the volume of the reactants / the depth of the reaction mixture [1 mark].
- 2.4 It would be more accurate to measure the volume of gas produced [1 mark] as this method less subjective [1 mark].
- 3.1 E.g. a gas syringe / a measuring cylinder inverted in a bowl of water [1 mark].




[2 marks for all points plotted correctly, or 1 mark for at least 5 points plotted correctly, 1 mark for line of best fit.]

- 3.3 Any value between 210-240 s [1 mark]
- When no more gas is produced, the reaction has stopped.
- 3.4 E.g. Mean rate of reaction = $\frac{\text{amount of product formed}}{\text{time for reaction to stop}}$
 $= \frac{18.0}{240} = 0.075 \text{ cm}^3/\text{s}$

[2 marks for correct answer between 0.075-0.086 cm³/s, otherwise 1 mark for correct equation]

If you got the wrong answer in 3.3, but used it correctly here as the change in y, you still get all the marks.

- 3.5 E.g. repeat the experiment using the same method [1 mark] and check that the results are similar [1 mark].




Gradient = $\frac{\text{change in } y}{\text{change in } x} = \frac{0.97}{153} = 0.0063 \text{ mol/s}$

(allow between 0.0053 mol/s and 0.0073 mol/s)

[4 marks for correct answer, otherwise 1 mark for correctly drawn tangent to curve at 50 s, 1 mark for answer to 2 s.f., 1 mark for correct units]

4.2



$$\text{Gradient} = \frac{\text{change in } y}{\text{change in } x} = \frac{0.45}{340} = 0.0013 \text{ mol/s}$$

(allow between 0.0008 mol/s and 0.0018 mol/s)

[4 marks for correct answer, otherwise 1 mark for correctly drawn tangent to curve at 200 s, 1 mark for answer to 2 s.f., 1 mark for correct units]

- 4.3 The rate decreases [1 mark]. This is because, as the amount of reactant A falls, so does its concentration and so the frequency of collisions between the reactant particles decreases [1 mark].

Page 71 — Reversible Reactions

- 1.1 That the reaction is reversible / can go both ways [1 mark].
 1.2 At equilibrium, the rate of the forward reaction is equal to the rate of the backwards reaction [1 mark].
 2.1 It will be exothermic [1 mark]. The same amount of energy will be released in the reverse reaction as is taken in by the forward reaction [1 mark].
 2.2 The system has reached equilibrium [1 mark]. This mixture contains both blue copper(II) ions and the yellow copper compound, so the colours mix to form green [1 mark].
 2.3 E.g. by changing the temperature / by changing the concentration of one of the reactants [2 marks — 1 mark for each correct answer].

Pages 72-73 — Le Chatelier's Principle

Warm-up

more reactants
 more reactants
 more products

- 1.1 If you change the conditions of a reversible reaction at equilibrium, the system will try to counteract that change [1 mark].
 1.2 E.g. the temperature / the concentration of the reactants [2 marks — 1 mark for each correct answer]
 2.1 At higher temperatures there will be more ICl and less ICl₃ / the equilibrium will shift to the left [1 mark]. This is because the reverse reaction is endothermic so opposes the increase in temperature [1 mark].
 2.2 There would be more ICl₃ and less ICl [1 mark] because the increase in pressure [1 mark] causes the equilibrium position to move to the side with the fewest molecules of gas [1 mark].
 3.1 At higher temperature there's more product (brown NO₂) in the equilibrium mixture [1 mark]. This suggests that the equilibrium has moved to the right/forward direction [1 mark], so the forward reaction is endothermic [1 mark].

From Le Chatelier's principle, you know that increasing the temperature will favour the endothermic reaction as the equilibrium tries to oppose the change. So the forward reaction must be endothermic, as there's more NO₂ in the equilibrium mixture at higher temperatures.

- 3.2 The mixture would go a darker brown [1 mark], as the decrease in pressure causes the equilibrium to move to the side with the most molecules of gas [1 mark], meaning more NO₂ is formed [1 mark].
 4 Observation 1: Increasing amounts of red FeSCN²⁺ are formed, so the solution becomes a darker red [1 mark]. When equilibrium is reached, the amount of each substance stops changing, and so does the colour [1 mark].
 Observation 2: The concentration of Fe³⁺ initially increases, so the solution becomes more orangey [1 mark]. The equilibrium then shifts to make more FeSCN²⁺, so the solution becomes darker red in colour [1 mark].
 Observation 3: The concentration of FeSCN²⁺ initially increases, so the solution becomes darker red [1 mark]. The equilibrium then shifts to produce more reactants, so the solution becomes more orangey [1 mark].

Topic 7 — Organic Chemistry

Pages 74-75 — Hydrocarbons

Warm-up


Hydrocarbon	Not a hydrocarbon
propane	butanoic acid
ethene	CH ₃ CH ₂ Cl
C ₂ H ₆	hydrochloric acid
C ₂ H ₄	

- 1.1 A compound that is formed from hydrogen and carbon atoms only [1 mark].
 1.2 butane, propane, ethane, methane [1 mark]
 1.3 C_nH_{2n+2} [1 mark]
 1.4 hydrocarbon + oxygen → carbon dioxide + water [1 mark]
 1.5 oxidised [1 mark]
 2.1 B [1 mark]
 2.2 B, D, and E [1 mark]. They have the general formula C_nH_{2n+2} [1 mark].
 2.3 E [1 mark]. Boiling point increases with increasing molecular size/number of carbons [1 mark].
 3.1 Diesel will be more viscous than petrol [1 mark]. The higher boiling point of diesel means it contains larger molecules/ molecules with longer chains [1 mark].
 3.2 Petrol [1 mark]. The lower boiling point of petrol means it contains smaller molecules/molecules with shorter chains [1 mark].
 3.3 C₂₀H₄₂ [1 mark]
 3.4 2C₈H₁₈ + 25O₂ → 16CO₂ + 18H₂O [1 mark for correct formulas of products, 1 mark for balancing]
 Any correct balance of the equation is correct, e.g. C₈H₁₈ + 12½O₂ → 8CO₂ + 9H₂O.

Page 76 — Fractional Distillation

- 1.1 The remains of ancient organisms/plankton [1 mark].
 1.2 A resource which is being used quicker than it is being replaced so will run out eventually [1 mark].
 1.3 alkanes [1 mark]
 2.1 boiling point [1 mark]
 2.2 The fractionating column is hot at the bottom and cool at the top [1 mark]. So longer hydrocarbons, which have higher boiling points, will condense and be drained off near the bottom [1 mark]. Meanwhile, shorter hydrocarbons, with lower boiling points, will condense and be drained off further up the column [1 mark].
 2.3 They contain similar numbers of carbon atoms / they have a similar chain length [1 mark].

Pages 77-78 — Uses and Cracking of Crude Oil

- 1.1 Any two from: e.g. solvents / lubricants / polymers / detergents [2 marks — 1 mark for each correct answer]
- 1.2 cracking [1 mark]
- 1.3 E.g. shorter chain hydrocarbons are more useful/can be used for more applications [1 mark].
- 2.1 thermal decomposition / endothermic [1 mark]
- 2.2 Hydrocarbons are vaporised / heated to form gases [1 mark].
The vapours are then passed over a hot catalyst / the vapours are mixed with steam and heated to very high temperatures [1 mark].
- 2.3 E.g. $C_{10}H_{22} \rightarrow C_7H_{16} + C_3H_6$ [1 mark]
Cracking equations must always be balanced and have a shorter alkane and an alkene on the right-hand side.
- 3.1 C_7H_{16} [1 mark]
- 3.2 
[1 mark for correct number of carbons, 1 mark for correct displayed formula]
- 3.3 E.g. to produce polymers / as a starting material for other chemicals [1 mark].
- 4 How to grade your answer:
Level 0: Nothing written worth of credit [No marks].
Level 1: Basic outline of how some fractions are processed but lacking detail. Some mention of the uses of cracking products [1 to 2 marks].
Level 2: Reason for cracking explained and some detail given about the process. The uses of cracking products are covered in detail [3 to 4 marks].
Level 3: Reasons for cracking and the process of cracking are explained in detail, including an accurate balanced symbol or word equation. Examples given of the uses of the products of cracking [5 to 6 marks].

Here are some points your answer may include:

Reasons for cracking

There is a higher demand for short chain hydrocarbons as these make good fuels.

Long chain hydrocarbons are less useful than short chain hydrocarbons, so there is less demand for them.


Cracking helps the supply of short chain hydrocarbons to meet the demand.

Cracking process

The long chain hydrocarbons are heated and vaporised. The vapours are passed over a hot catalyst / mixed with steam and heated to a high temperature so that they thermally decompose.

Any relevant word equation: e.g. decane \rightarrow octane + ethene


Any relevant balanced equation:

**Uses of cracking products**

The products of cracking are useful as fuels.

Alkenes are used as a starting material when making lots of other compounds and can be used to make polymers.

Page 79 — Alkenes

- 1.1 A hydrocarbon with a double carbon-carbon bond [1 mark].
- 1.2 C_nH_{2n} [1 mark]
- 1.3 
[1 mark]
- 2.1 **B and E** [1 mark]
- 2.2 C_6H_{12} [1 mark]


- 2.3 **B** undergoes incomplete combustion in low amounts of oxygen [1 mark].
Any balanced incomplete combustion equation:
e.g. $2C_2H_4 + 5O_2 \rightarrow 2CO_2 + 2CO + 4H_2O$ [1 mark for correct formulas of reactants and products, 1 mark for balancing]

Pages 80-81 — Reactions of Alkenes


Warm-up

Alkenes generally react via **addition** reactions to form a variety of compounds. Alkenes can react with steam to form **alcohols**. For example, **ethene** can be mixed with steam and passed over a catalyst to form ethanol.



- 1.1 $C=C$ / carbon-carbon double bond [1 mark]
- 1.2 They have the same functional group [1 mark].
- 1.3 The double bond opens up to leave a single bond [1 mark].
A new atom is added to each of the $C=C$ carbons [1 mark].
- 2.1 C_3H_6 [1 mark]
- 2.2 a catalyst / nickel [1 mark]
- 2.3 Add bromine water [1 mark]. With propene it will change from orange to colourless [1 mark]. Propane will not react and there will be no colour change [1 mark].





- 3.2 The product of the reaction is saturated [1 mark]. It doesn't contain any carbon-carbon double bonds [1 mark].




You still get the mark if you have the chlorine and iodine atoms swapped around.

**Page 82 — Addition Polymers**

- 1.1 ethene [1 mark]
- 1.2 addition polymerisation [1 mark]
- 1.3 carbon-carbon double bond / $C=C$ [1 mark]
- 2.1 
[1 mark]
- 2.2 
[1 mark]
- 2.3 chloroethene [1 mark]

Pages 83-84 — Alcohols

- 1.1 $-OH$ [1 mark]
- 1.2 Any two from: e.g. fuels / solvents / in alcoholic drinks [2 marks — 1 mark for each correct answer].
- 1.3 Methanol dissolves. The indicator remains green [1 mark].
- 1.4 CO_2 and H_2O [1 mark]
- 2.1 Ethanoic acid [1 mark], a carboxylic acid [1 mark].
- 2.2 Yeast [1 mark]. The reaction conditions are a temperature of approximately $37^\circ C$ [1 mark], absence of oxygen / anaerobic conditions [1 mark] and a slightly acidic pH [1 mark].



3.2 Methanol only has one carbon atom whereas butanol has four/methanol has the formula CH_3OH whereas butanol has the formula of $\text{C}_4\text{H}_9\text{OH}$ [1 mark]. Both methanol and butanol contain an $-\text{OH}$ group [1 mark].

3.3 They both contain the same functional ($-\text{OH}$) group [1 mark].

3.4 hydrogen [1 mark]

4.1 $\text{C}_2\text{H}_4(\text{OH})_2/\text{C}_2\text{H}_6\text{O}_2$ [1 mark]

4.2 E.g. react it with sodium [1 mark].

4.3 $2\text{C}_2\text{H}_4(\text{OH})_2 + 5\text{O}_2 \rightarrow 4\text{CO}_2 + 6\text{H}_2\text{O}$ [1 mark for the formulas of the reactants and products, 1 mark for balancing]

Any correct balance of the equation is correct, e.g. $\text{C}_2\text{H}_4(\text{OH})_2 + 2\frac{1}{2}\text{O}_2 \rightarrow 2\text{CO}_2 + 3\text{H}_2\text{O}$.

4.4 7/neutral [1 mark]

Pages 85-86 — Carboxylic Acids

1.1 $-\text{COOH}$ [1 mark]

1.2 Methanoic acid would dissolve [1 mark] and the Universal indicator would change to red/orange [1 mark].

1.3 carbon dioxide/ CO_2 [1 mark]

2.1 butanoic acid [1 mark], $\text{C}_3\text{H}_7\text{COOH}/\text{C}_4\text{H}_8\text{O}_2$ [1 mark]

2.2 It does not ionise completely when dissolved in water [1 mark].

2.3 An acid catalyst [1 mark].

2.4 ester [1 mark]

3.1 Accept anywhere between 2 and 6 [1 mark].

3.2 magnesium carbonate/ MgCO_3 [1 mark]

3.3 ethanol/ $\text{C}_2\text{H}_5\text{OH}$ [1 mark]

4.1 Na_2CO_3 [1 mark]

4.2 H_2O [1 mark]

4.3 **D** [1 mark]. Propanoic acid is a weak acid [1 mark] and therefore weakly ionised [1 mark].

4.4 Carbon dioxide/ CO_2 [1 mark] and water/ H_2O [1 mark].


Page 87 — Condensation Polymers

1.1 A small molecule is lost when condensation polymers are formed [1 mark].

1.2 two [1 mark]

1.3 E.g. carboxylic acids [1 mark] and alcohols / amines [1 mark].

2.1 H_2O [1 mark]



[1 mark for correct ester link connecting monomers, 1 mark for rest of the structure being correct]

2.3 No [1 mark]. As two different functional groups that react with each other are needed [1 mark].

Page 88 — Naturally Occurring Polymers

1.1 two [1 mark]

1.2 Any one from: carboxylic acid / amine [1 mark]

1.3 condensation polymerisation [1 mark]

1.4 water [1 mark]

2.1 sugars [1 mark]

2.2 e.g. cellulose [1 mark]

2.3 DNA contains genetic instructions [1 mark] for the operation and functioning of living organisms and viruses [1 mark].

2.4 Two polymer chains [1 mark] made from four different nucleotide monomers [1 mark] linked together by cross links to give a double helix structure [1 mark].


Topic 8 — Chemical Analysis

Page 89 — Purity and Formulations

- 1.1 A single element or compound not mixed with any other substance [1 mark].
- 1.2 Sample A [1 mark]. The purer the substance, the smaller the range of the melting point / purer substances melt at higher temperatures than impure substances [1 mark].
- 1.3 Sample A [1 mark].
- 2.1 It is a mixture that has been designed to have a precise purpose [1 mark]. Each of the components is present in a measured quantity [1 mark] and contributes to the properties of the formulation [1 mark].
- 2.2 By making sure each component in the mixture is always present in exactly the same quantity [1 mark].
- 2.3 Any one from: e.g. medicines / cleaning products / fuels / cosmetics / fertilisers / metal alloys [1 mark].

Pages 90-91 — Paper Chromatography

Warm-up



- 1.1 $E: R_f = \frac{\text{distance travelled by substance}}{\text{distance travelled by solvent}} = \frac{3.6}{9.5} = 0.38$
[2 marks for correct answer, otherwise 1 mark for using correct equation to calculate R_f]
- $F: R_f = \frac{\text{distance travelled by substance}}{\text{distance travelled by solvent}} = \frac{8.0}{9.5} = 0.84$
[2 marks for correct answer, otherwise 1 mark for using correct equation to calculate R_f]
- 1.2 E.g. they're distributed differently between the mobile phase and the stationary phase [1 mark].
- 1.3 They're all pure substances [1 mark].
- 1.4 **D** and **E** [1 mark].
- 2.1 E.g. to stop any solvent evaporating [1 mark].
- 2.2 **A** spends more time in the mobile phase compared to the stationary phase than **B** does [1 mark].
- 2.3 **B** and **C** [1 mark].
- 2.4 The student is incorrect [1 mark]. Substances have different R_f values in different solvents as the attraction between the substance and solvent changes [1 mark].
- 2.5 It suggest that there are at least 3 substances in **W** [1 mark].
- 2.6 There were only two spots in the chromatogram shown because two of the substances in **W** are similarly distributed between the mobile phase/water and stationary phase / they had similar R_f values [1 mark].

Pages 92-93 — Tests for Gases and Anions

- 1.1 A burning splint which results in a popping noise [1 mark].
- 1.2 oxygen/ O_2 [1 mark]
- 1.3 Bubble the gas through an aqueous solution of calcium hydroxide/lime water [1 mark]. The lime water turns milky/cloudy [1 mark].
- 1.4 Damp litmus paper [1 mark] put into the gas. If chlorine gas is present, the paper is bleached and turns white [1 mark].
- 2.1 Potassium, K^+ [1 mark]
- 2.2 Clean a platinum wire loop [1 mark] by dipping it in some dilute HCl and then placing it in a blue flame from a Bunsen burner until it burns without colour [1 mark]. Dip the loop into the sample you want to test and put it back into the flame [1 mark]. Record the colour of the flame [1 mark].
- 2.3 yellow [1 mark]
- 3.1 sulfate/ SO_4^{2-} [1 mark]

- 3.2 $\text{Ba}^{2+}_{(\text{aq})} + \text{SO}_4^{2-}_{(\text{aq})} \rightarrow \text{BaSO}_{4(\text{s})}$ [1 mark for balanced equation, 1 mark for state symbols]
- 3.3 copper(II)/ Cu^{2+} [1 mark]
- 3.4 copper sulfate/ CuSO_4 [1 mark]
- 4.1 Substance P: CaI_2 /calcium iodide [1 mark for correct anion, 1 mark for correct cation]
Substance R: LiBr/lithium bromide [1 mark for correct anion, 1 mark for correct cation]
- 4.2 The anion cannot be identified [1 mark].
- 4.3 D: white precipitate [1 mark]
E: green [1 mark]
- 4.4 The flame colours of some ions may be hidden by / mixed with the colour of others [1 mark].

Page 94 — Flame Emission Spectroscopy

- 1.1 Any two from: e.g. faster / more sensitive / more accurate [2 marks — 1 mark for each correct answer]
- 1.2 The identity [1 mark] and the concentration of metal ions in solution [1 mark].
- 1.3 line spectra [1 mark]
- 2.1 A sample is placed in a flame and as the ions in the sample heat up they transfer energy as light [1 mark]. This light passes through a spectroscope and produces a line spectrum specific to that ion [1 mark].
- 2.2 Metal A and metal C [1 mark].

Topic 9 — Chemistry of the Atmosphere

Pages 95-96 — The Evolution of the Atmosphere

Warm-up

- 1 False
- 2 True
- 3 True
- 4 False
- 1.1 One-fifth oxygen and four-fifths nitrogen [1 mark].
- 1.2 Any two from: e.g. carbon dioxide / water vapour / named noble gas [2 marks — 1 mark for each correct answer]
- 1.3 By algae and plants photosynthesising [1 mark].
- 1.4 By volcanic activity [1 mark].
- 1.5 200 million years [1 mark]
- 2.1 E.g. photosynthesis by plants and algae / carbon dioxide dissolved in the oceans [1 mark].
- 2.2 From matter that is buried and compressed over millions of years [1 mark].
- 2.3 Coal: from thick plant deposits [1 mark].
Limestone: from calcium carbonate deposits from the shells and skeletons of marine organisms [1 mark].
- 3.1 E.g. the long timescale means there's a lack of evidence [1 mark].
- 3.2 $6\text{CO}_2 + 6\text{H}_2\text{O} \rightarrow \text{C}_6\text{H}_{12}\text{O}_6 + 6\text{O}_2$ [1 mark]
- 3.3 Oxygen is produced by photosynthesis [1 mark] and there are no plants or algae / there isn't any photosynthesis [1 mark] on Mars.
- 3.4 The fact that the red beds formed about 2 billion years ago suggests that before this time there wasn't enough oxygen in the air for iron oxide to form / from this time there was enough oxygen in the air for iron oxide to form [1 mark].

Pages 97-98 — Greenhouse Gases and Climate Change

- 1.1 Nitrogen [1 mark]
They help to keep Earth warm [1 mark].
- 1.3 Any two from: e.g. deforestation / burning fossil fuels / agriculture / producing waste [2 marks — 1 mark for each correct answer]
- 2.1 Greenhouse gases absorb long-wave (thermal) radiation [1 mark] reflected from Earth's surface [1 mark]. They then reradiate this thermal radiation in all directions, including back towards Earth, helping to warm the atmosphere [1 mark].

- 2.2 E.g. flooding [1 mark] and coastal erosion [1 mark].
- 2.3 Any one from: e.g. changes in rainfall patterns / the ability of certain regions to produce food might be affected / the frequency/severity of storms might increase / the distribution of wild species might change [1 mark].
- 3 How to grade your answer:
Level 0: There is no relevant information [No marks].
Level 1: Unstructured and no logic. The trends in the variables are described but reasons are not given [1 to 2 marks].
Level 2: Some structure and logic but lacking clarity. The trends in the variables are described and there is some explanation of how the increase in carbon dioxide may have come about and how this might be linked to temperature [3 to 4 marks].
Level 3: Clear, logical answer. The trends in the variables are described and there is a clear explanation of how the increase in carbon dioxide may have come about and how this may be linked to temperature [5 to 6 marks].

Here are some points your answer may include:

The graph shows an increase in carbon dioxide levels in the atmosphere between 1960 and 2015.

The increase in carbon dioxide levels is likely to be due to human activities which release carbon dioxide into the atmosphere.

These activities include increased burning of fossil fuels, increased deforestation and increased waste production.

The graph shows that the increase in carbon dioxide appears to correlate with an increase in global temperatures.

The increase in global temperatures is likely to be due to the increase in carbon dioxide in the atmosphere, as carbon dioxide is a greenhouse gas so helps to keep Earth warm.

4.1 The global warming potential for methane is significantly greater than for carbon dioxide [1 mark].

4.2 It has a very high global warming potential compared to other gases [1 mark] and stays in the atmosphere for a long time [1 mark].

Page 99 — Carbon Footprints

- 1.1 A measure of the amount of carbon dioxide and other greenhouse gases [1 mark] released over the full life cycle of something [1 mark].
- 1.2 E.g. using renewable or nuclear energy sources [1 mark] and using more energy efficient appliances [1 mark].
- 1.3 E.g. lack of education / reluctance to change their lifestyle / cost of changing lifestyle [1 mark].
- 2.1 Any two from, e.g: specialist equipment is needed to capture the carbon dioxide / it's expensive to capture and store the carbon dioxide / it could be difficult to find suitable places to store the carbon dioxide [2 marks — 1 mark for each correct answer].
- 2.2 E.g. governments could tax companies based on the amount of greenhouse gases they emit [1 mark]. They could also put a cap on the emissions produced by a company [1 mark]. Governments might be reluctant to impose these methods if they think it will affect economic growth / could impact on people's well-being [1 mark], especially if other countries aren't using these methods either / the country is still developing [1 mark].

Page 100 — Air Pollution

- 1.1 Coal can contain sulfur impurities [1 mark]
- 1.2 Acid rain: sulfur dioxide / nitrogen oxides/nitrogen monoxide/nitrogen dioxide/dinitrogen monoxide [1 mark]
Global dimming: e.g. (carbon) particulates [1 mark]
- 1.3 Any two from: e.g. damage to plants / buildings / statues / corrodes metals [2 marks — 1 mark for each correct answer].

- 2.1 The reaction of nitrogen and oxygen from the air [1 mark] at the high temperatures produced by combustion [1 mark].
- 2.2 Nitrogen oxides cause respiratory problems [1 mark] and contribute to acid rain [1 mark].
- 2.3 E.g. they can cause respiratory problems [1 mark].
- 2.4 Carbon monoxide [1 mark]. It is colourless and odourless [1 mark].

Topic 10 — Using Resources

Page 101 — Ceramics, Polymers and Composites

- 1.1 limestone, sand, sodium carbonate [3 marks — 1 mark for each correct answer]
- 1.2 Because borosilicate glass has a higher melting point than soda-lime glass [1 mark].
- 1.3 Wet clay is shaped [1 mark] then fired at a high temperature [1 mark].
- 2.1 They are made at different temperatures/pressures [1 mark] with a different catalyst [1 mark].
- 2.2 thermosoftening [1 mark]
- 2.3 Poly(ethene) chains are entwined together with weak forces between the chains [1 mark]. Polyester resin can form crosslinks between polymer chains [1 mark].
- 2.4 Composites consist of fibres/fragments of a material known as the reinforcement [1 mark] surrounded by a matrix/binder [1 mark].

Page 102 — Properties of Materials

- 1.1 Bronze — Copper and tin [1 mark]
Steel — Iron and carbon [1 mark]
Brass — Copper and zinc [1 mark]
- 1.2 Any one from: e.g. water taps / door fittings [1 mark]
- 2.1 strength increases [1 mark]
- 2.2 high carbon steel [1 mark]
- 2.3 Aluminium is much less dense than the other metals [1 mark].

Page 103 — Corrosion

- 1.1 iron + water + oxygen → hydrated iron(III) oxide [1 mark]
- 1.2 Galvanising is coating iron with zinc [1 mark]. Zinc is more reactive than iron [1 mark] so, unlike other methods of barrier protection, even if the coating is scratched it will still prevent the iron from rusting as it will oxidise before iron does [1 mark].
- 1.3 Any three from: e.g. painting / coating with grease / electroplating / using a barrier / sacrificial protection [3 marks — 1 mark for each correct answer]
- 2.1 The reaction of a material with substances in its environment so it is gradually destroyed [1 mark].
- 2.2 The oxide that forms when aluminium reacts in the air forms a protective layer over the surface of the metal [1 mark], preventing chemicals reaching the rest of the metal and reacting further [1 mark].
- 2.3 Oxygen and water react with the magnesium instead of the steel [1 mark].

Page 104 — Finite and Renewable Resources

- 1.1 Coal [1 mark]. It does not form fast enough to be considered replaceable [1 mark].
- 1.2 A resource that reforms at a similar rate to, or faster, than humans can use it [1 mark].
- 2.1 E.g. the development of fertilisers has meant higher yields of crops [1 mark].

- 2.2 Any one from: e.g. synthetic rubber has replaced natural rubber / poly(ester) has replaced cotton in clothes / bricks are used instead of timber in construction [1 mark].
- 3 Any one advantage from: e.g. allows useful products to be made / provides jobs / brings money into the area [1 mark]. Any one disadvantage: e.g. uses large amounts of energy / scars the landscape / produces lots of waste / destroys habitats [1 mark].

Pages 105-106 — Reuse and Recycling

- 1.1 An approach to development that takes account of the needs of present society [1 mark] while not damaging the lives of future generations [1 mark].
- 1.2 E.g. chemists can develop and adapt processes that use less resources/do less damage to the environment [1 mark]. For example, chemists have developed catalysts that reduce the amount of energy required for industrial processes [1 mark].
- 2.1 The raw materials for the jute bag are more sustainable [1 mark] as plant fibres are a renewable resource, whilst crude oil is a finite resource [1 mark].
- 2.2 The production of the poly(ethene) bag is more sustainable [1 mark] as it needs less energy to be produced from its raw materials than the jute bag [1 mark].
- 2.3 The jute bag can be reused and the poly(ethene) bag can be recycled, improving both their sustainability [1 mark]. However, the jute bag is more sustainable if the bags are disposed of in landfill [1 mark], as it is biodegradable, whilst the poly(ethene) bag isn't [1 mark].
- 3.1 Any two from: e.g. often uses less energy / conserves the amount of raw materials on Earth / cuts down on waste sent to landfill [2 marks — 1 mark for each correct answer].
- 3.2 Any one from: e.g. glass / metal [1 mark]
E.g. glass is crushed and melted down to form other glass products/other purpose / metal is melted and cast into the shape of a new product [1 mark].
- 3.3 reusing [1 mark]
- 4.1 Plants are grown on soil containing copper compounds [1 mark], so as they grow, copper builds up in their leaves [1 mark]. The plants are burned [1 mark]. The resulting ash contains the copper compounds [1 mark].
- 4.2 By electrolysis of a solution containing the copper compounds [1 mark] or by displacement using scrap iron [1 mark].
- 4.3 Copper is a finite resource [1 mark] and will eventually run out [1 mark]. Recycling copper makes it more sustainable [1 mark].

Page 107 — Life Cycle Assessments

- Warm-up
- Getting the Raw Materials — Coal being mined from the ground.
- Manufacturing and Packaging — Books being made from wood pulp.
- Using the Product — A car using fuel while driving.
- Product Disposal — Plastic bags going on to landfill.
- 1.1 Any two from: e.g. if a product is disposed of in landfill sites, it will take up space / may pollute land/water / energy is used to transport waste to landfill / pollution can be caused by incineration [2 marks — 1 mark for each correct answer].
- 1.2 Any one from: e.g. energy / water / some natural resources / certain types of waste [1 mark]
- 1.3 They can be subjective / they are difficult to measure [1 mark].
- 1.4 No [1 mark]. Some elements of the LCA are not objective/require the assessors to make value judgements/cannot be quantified reliably [1 mark], therefore different people are likely to make a different judgement/estimate [1 mark].
- 1.5 Selective LCAs could be written so they only show elements that support a company's claims / they could be biased [1 mark] in order to give them positive advertising [1 mark].

Pages 108-109 — Potable Water

Warm-up

- 1 False
2 True
3 False
- 1.1 pure water [1 mark]
1.2 e.g. from the ground / lakes / rivers [1 mark].
1.3 passing water through filter beds — solid waste [1 mark]
sterilisation — microbes [1 mark]
1.4 E.g. chlorine, ozone, ultraviolet light [3 marks — 1 mark for each correct answer].
- 2.1 A: Bunsen burner [1 mark]
B: round bottom flask [1 mark]
C: thermometer [1 mark]
D: condenser [1 mark]
- 2.2 Pour the salt water into the flask and secure it on top of a tripod [1 mark]. Connect the condenser to a supply of cold water [1 mark] that goes in at the bottom and out at the top [1 mark]. Heat the flask and allow the water to boil [1 mark]. Collect the water running out of the condenser in a beaker [1 mark].
- 2.3 Reverse osmosis / a method which uses membranes [1 mark]
2.4 Desalination requires a lot of energy compared to the filtration and sterilisation of fresh water [1 mark]. Since the UK has a plentiful supply of fresh water there is no need to use desalination processes [1 mark].

Page 110 — Waste Water Treatment

- 1.1 organic matter, harmful microbes [2 marks — 1 mark for each correct answer]
1.2 It may contain harmful chemicals which need to be removed [1 mark].
2.1 To remove grit [1 mark] and large bits of material/twigs/plastic bags [1 mark].
2.2 Substance A: sludge [1 mark]
Substance B: effluent [1 mark]
2.3 anaerobic digestion [1 mark]

Page 111 — The Haber Process



- 1.1 Nitrogen, Hydrogen [1 mark for both]
1.2 ammonia [1 mark]
1.3 Because ammonia is used to make fertilisers [1 mark].
2.1 Low temperature [1 mark]. Low temperatures cause the position of equilibrium to shift in favour of the exothermic, forward reaction [1 mark] which means more ammonia is produced/you get a higher yield of ammonia [1 mark].
2.2 A higher temperature is used to get a reasonable rate of reaction [1 mark].
2.3 High pressure causes the yield to increase [1 mark]. There are fewer moles of gas in the products than reactants/ on the right hand side of the equation than on the left hand side [1 mark]. Since high pressure favours the production of fewer moles of gas, the position of equilibrium would move right/to the product side as pressure is increased [1 mark].
2.4 Any one from: e.g. safety / cost / rate [1 mark for any correct answer].

Page 112 — NPK Fertilisers

- 1.1 potassium chloride, potassium sulfate [1 mark for each correct answer]
1.2 They are mined [1 mark].
2.1 calcium nitrate [1 mark]
2.2 Phosphate rock + sulfuric acid: calcium phosphate [1 mark] and calcium sulfate [1 mark].
Phosphate rock + phosphoric acid: calcium phosphate [1 mark].
2.3 The reaction is carried out at lower concentrations in the lab so that it's safer for the person carrying it out [1 mark]. Crystallisation isn't used in industry as it's very slow [1 mark].

Mixed Questions

Pages 113-124 — Mixed Questions

- 1.1
- 
- 2 marks if all three correct, otherwise 1 mark if 1 correct
- 1.2 Mixtures with a precise purpose [1 mark] that are made by following a formula / a recipe [1 mark].
2.1 Dissolve the rock salt in water and filter [1 mark].
2.2 It contains two elements/more than one element in fixed proportions [1 mark] held together by chemical bonds [1 mark].
2.3 ionic [1 mark]
3.1 Group: 6 [1 mark]
Explanation: There are 6 electrons in the outer shell [1 mark].
3.2 2- ions [1 mark], as oxygen atoms need to gain two electrons to get a full outer shell [1 mark].
3.3 Oxidation [1 mark]
- 4.1
- 
- 1 mark for shared pair of electrons, 1 mark for six further electrons in the outer shell of each chlorine atom
- 4.2 E.g. atoms with the same number of protons / of the same element / with the same atomic number [1 mark] with different numbers of neutrons / different mass numbers [1 mark].
4.3 Hold a piece of damp litmus paper in the unknown gas [1 mark]. It will be bleached white in the presence of chlorine [1 mark].
4.4 Chlorine is more reactive than iodine [1 mark], so would displace iodine from sodium iodide solution / the solution would go from colourless to brown [1 mark].
5.1 endothermic [1 mark]
5.2 higher [1 mark]
5.3 It takes more energy to break the bonds in the reactants than is released when the bonds in the products form [1 mark], so overall energy is taken in from the surroundings [1 mark].
5.4 E.g. in a sports injury pack [1 mark].
6.1 alkanes [1 mark]
6.2 (fractional) distillation [1 mark]
6.3 cracking [1 mark]
6.4 Decane [1 mark], because the molecules are bigger [1 mark], so will have stronger intermolecular forces / more energy is needed to break the forces between the molecules [1 mark].
6.5 $C_5H_{12} + 8O_2 \rightarrow 5CO_2 + 6H_2O$ [1 mark for correct reactants and products, 1 mark for balancing]
7.1 The electrons in the outer shell [1 mark] of the metal atoms are delocalised [1 mark]. There is strong electrostatic attraction between the positive metal ions and the shared negative electrons [1 mark].
7.2 Iron: solid [1 mark]. Silver: liquid [1 mark]
7.3 Iron [1 mark], because it has a higher melting/boiling point [1 mark], so more energy is needed to break the bonds [1 mark].
8.1 The reactant that is used up first / limits the amount of product that's formed [1 mark].
8.2 $M_r(LiOH) = A_r(Li) + A_r(O) + A_r(H) = 7 + 16 + 1 = 24$ [1 mark]
8.3 Number of moles = mass ÷ molar mass = $1.75 \div 7 = 0.25 \text{ mol}$ [2 marks for correct answer, otherwise 1 mark for using the correct equation to calculate moles]

- 8.4 From the reaction equation, 0.50 mol Li forms 0.50 mol LiOH.
Mass of LiOH = number of moles \times molar mass = $0.50 \times 24 = 12 \text{ g}$ [3 marks for correct answer, otherwise 1 mark for number of moles of LiOH produced, 1 mark for using the correct equation to calculate mass]
- 9.1 It described atoms as having a tiny, positively charged nucleus at the centre [1 mark], surrounded by a cloud of electrons [1 mark].
- 9.2 Atoms consist of a small nucleus [1 mark] which contains the protons and neutrons [1 mark]. The electrons orbit the nucleus in fixed energy levels/shells [1 mark].
- 10.1 The particles in a gas expand to fill any container they're in [1 mark]. So the particles of carbon dioxide formed will expand out of the unsealed reaction vessel [1 mark], causing the mass of substance inside the reaction vessel to decrease [1 mark].
- 10.2 E.g. add a set volume and concentration of hydrochloric acid to the reaction vessel [1 mark]. Add a set volume and concentration of sodium carbonate solution [1 mark], connect the reaction flask to a gas syringe [1 mark] and start the stop-watch [1 mark]. Record the volume of gas collected at regular intervals until the reaction is finished [1 mark]. Repeat the experiment, keeping everything the same except for the concentration of acid [1 mark].
- 10.3 Change in volume = 12.0 cm^3
Mean rate of reaction = $\frac{\text{amount of product formed}}{\text{time}} = \frac{12.0}{30} = 0.40 \text{ cm}^3/\text{s}$ [2 marks for correct answer, otherwise 1 mark for using the correct equation to calculate rate]
- 11.1 Any two from: e.g. it dissolved in oceans / photosynthesis / trapped in rocks and fossil fuels [2 marks — 1 mark for each correct answer]
- 11.2 E.g. methane [1 mark]. It is increasing due to more agriculture / waste production [1 mark].
- 11.3 How to grade your answer:
Level 0: There is no relevant information. [No marks]
Level 1: There are a few examples of other pollutant gases, but little discussion of how they are made or what their impacts could be. [1 to 2 marks]
Level 2: There are a number of examples of other pollutant gases, with some discussion of how they are made and what their impacts could be. [3 to 4 marks]
Level 3: There are a number of examples of other pollutant gases, with a detailed discussion of how they are made and what their impacts could be. [5 to 6 marks]
- Here are some points your answer may include:
Other pollutant gases include carbon monoxide, sulfur dioxide and nitrogen oxides.
Carbon monoxide is produced when fuels undergo incomplete combustion.
Carbon monoxide can cause fainting, coma or even death.
Sulfur dioxide is produced when fuels that contain sulfur impurities are burned.
Sulfur dioxide can mix with water in clouds to produce sulfuric acid, so cause acid rain.
Sulfur dioxide can cause respiratory problems.
Nitrogen oxides are produced when nitrogen and oxygen from the air react/combine due to the heat of burning.
Nitrogen oxides can mix with water in clouds to produce nitric acid, so cause acid rain.
Nitrogen oxides can cause respiratory problems.
- 12.1 Iron, tin, copper. [2 marks if all correct, or 1 mark if 1 correct]
- 12.2 $2\text{AgCl}_{(\text{aq})} + \text{Cu}_{(\text{s})} \rightarrow \text{CuCl}_{2(\text{aq})} + 2\text{Ag}_{(\text{s})}$ [1 mark for correct equation, 1 mark for balancing, 1 mark for state symbols]
- 13.1 Copper is lower in the reactivity series/less reactive than carbon [1 mark], so can be extracted by reduction using carbon [1 mark].
- 13.2 Bacteria are used to convert copper compounds in the ore into soluble copper compounds [1 mark]. This produces a leachate that contains copper ions [1 mark] which can be extracted by electrolysis/displacement with iron [1 mark].
- 13.3 The atoms in copper form layers which slide over each other, so it can be drawn out into wires [1 mark]. Copper contains delocalised electrons which are free to move and carry an electric current [1 mark].
- 13.4 The tin atoms in bronze distort the structure of the copper [1 mark]. This means the layers can no longer slide over each other [1 mark], so bronze is harder than copper [1 mark].
- 14.1 Polymer A has weak forces between the chains [1 mark]. Polymer B has cross links between the chains [1 mark].
- 14.2 Polymer B [1 mark] as it's rigid, so would keep the shape of the mug [1 mark] and it wouldn't be softened by the hot drinks [1 mark].
- 15.1
$$\begin{array}{c} \text{H} \quad \text{H} \\ | \quad | \\ \text{H}-\text{C}-\text{C}-\text{O}-\text{H} \\ | \quad | \\ \text{H} \quad \text{H} \end{array}$$
 [1 mark for correct number of each atom, 1 mark for atoms joined up correctly]
- 15.2 Ethanol can be made by reacting ethene with steam [1 mark] in the presence of a catalyst [1 mark]. It can also be made by fermenting sugars with yeast [1 mark] at around 37°C [1 mark] and slightly acidic conditions [1 mark] in the absence of oxygen [1 mark].
- 16.1 Add a few drops of sodium hydroxide to a sample of the solution [1 mark]. If iron(II) ions are present, a green precipitate should form [1 mark].
 $\text{FeSO}_{4(\text{aq})} + 2\text{NaOH}_{(\text{aq})} \rightarrow \text{Fe}(\text{OH})_{2(\text{s})} + \text{Na}_2\text{SO}_{4(\text{aq})}$ [1 mark for correct equation, 1 mark for balancing, 1 mark for state symbols]
- 16.2 Add some barium chloride [1 mark] to the solution in the presence of hydrochloric acid [1 mark]. If sulfate ions are present, a white precipitate should form [1 mark].
 $\text{BaCl}_{2(\text{aq})} + \text{FeSO}_{4(\text{aq})} \rightarrow \text{BaSO}_{4(\text{s})} + \text{FeCl}_{2(\text{aq})}$ [1 mark for balanced equation, 1 mark for state symbols]
- 16.3 Add iron(II) oxide to sulfuric acid until the reaction stops / the solid sinks to the bottom [1 mark]. Filter off the excess iron(II) oxide [1 mark]. Gently heat the iron(II) sulfate solution to evaporate some of the water and then leave to cool [1 mark]. Filter and dry the crystals that form [1 mark].
- 17.1 Number of moles of carbon = $\text{mass} \div A_r = 24 \div 12 = 2 \text{ mol}$
1 mol of carbon reacts to produce 2 mol of hydrogen gas, so 2 mol of carbon will react to produce $2 \times 2 = 4 \text{ mol}$ of hydrogen gas.
 M_r of $\text{H}_2 = 2 \times 1 = 2$
Mass = number of moles \times molar mass = $4 \times 2 = 8 \text{ g}$
[4 marks for correct answer, or 1 mark for correct number of moles of carbon, 1 mark for correct number of moles of hydrogen, 1 mark for correct M_r of hydrogen]
- 17.2 1 mole of gas occupies 24 dm^3 at room temperature and pressure, so 4 moles of gas occupies $4 \times 24 = 96 \text{ dm}^3$ [1 mark]
If you got the answer to 13.1 wrong you still get the mark here if you used your answer to 13.1 correctly in this part.
- 17.3 Percentage yield
$$= \frac{\text{Mass of product actually made}}{\text{Maximum theoretical mass of product}} \times 100 = \frac{4.8}{8} \times 100 = 60\%$$
 [2 marks for correct answer, otherwise 1 mark for using the correct equation to calculate percentage yield]
- 18.1 100% [1 mark]. The reaction only has one product [1 mark].
- 18.2 E.g. the reaction is reversible / some of the products will always turn back into reactants / there might be side reactions / some of the product may be lost as it's separated from the reaction mixture [1 mark].

- 18.3 A low temperature shifts the position of equilibrium in favour of the forward, exothermic reaction [1 mark]. This means there will be more product at equilibrium / the yield will be greater [1 mark]. However, a low temperature decreases the rate of reaction [1 mark]. So the temperature is a compromise in order to get a good yield at a reasonable rate [1 mark].
- 19.1 $H^+ + OH^- \rightarrow H_2O$ [1 mark]
- 19.2 sodium sulfate [1 mark]
- 19.3 Universal indicator doesn't have a sudden colour change at the endpoint [1 mark]. An indicator such as methyl orange / phenolphthalein / litmus [1 mark] should be used instead.
- 19.4 volume in $dm^3 = 20.35 \div 1000 = 0.02035 dm^3$
 moles = volume \times concentration = $0.02035 \times 0.10 = 0.0020 mol$ [3 marks for correct answer, otherwise 1 mark for volume in dm^3 , 1 mark for using the correct equation to calculate moles]
- 19.5 mean titre = $(20.05 + 19.95 + 20.00) \div 3 = 20.00 cm^3$ [2 marks for correct answer, otherwise 1 mark for ignoring rough titre]
- 19.6 volume of H_2SO_4 to react in $dm^3 = 20.0 \div 1000 = 0.0200 dm^3$
 moles of H_2SO_4 to react = $0.0200 \times 0.10 = 0.0020 mol$
 $H_2SO_4 + 2NaOH \rightarrow Na_2SO_4 + 2H_2O$
 1 mole of H_2SO_4 reacts with 2 moles of NaOH, so
 0.0020 mol of H_2SO_4 reacts with 0.0040 mol of NaOH
 Concentration of NaOH = number of moles \div volume
 = $0.0040 \div 0.025 = 0.16 mol/dm^3$ [5 marks for correct answer, otherwise 1 mark for number of moles of H_2SO_4 , 1 mark for number of moles of NaOH, 1 mark for balanced reaction equation, 1 mark for using the correct equation to calculate concentration]
- 20.1 Aluminium ore is mixed with cryolite and melted [1 mark]. An electric current is passed through the molten ore [1 mark]. At the cathode/negative electrode, Al^{3+} ions are reduced to aluminium metal [1 mark]: $Al^{3+} + 3e^- \rightarrow Al$ [1 mark]. At the anode/positive electrode, O^{2-} ions are oxidised to oxygen [1 mark]: $2O^{2-} \rightarrow O_2 + 4e^-$ [1 mark].
- 20.2 When aluminium corrodes it forms a protective layer of aluminium oxide [1 mark] that stops any further reaction taking place [1 mark].
- 20.3 Galvanising means covering iron with a layer of zinc [1 mark]. This acts as a protective barrier to keep out water and oxygen [1 mark]. If the layer is scratched, the zinc around the scratch reacts instead of the iron [1 mark].
- 21.1 Order: diamond, poly(propene), butane [1 mark].
 Explanation: Diamond has the highest melting point as you need to break the strong covalent bonds [1 mark]. Poly(propene) forms larger molecules than butane, so has stronger intermolecular forces [1 mark], which require more energy to break [1 mark].
- 21.2 Particles with a diameter between 1 nm and 100 nm / particles containing only a few hundred atoms [1 mark]. They have a very high surface area to volume ratio compared to bulk materials [1 mark].
- 21.3 E.g. they could be used for drug delivery [1 mark]. The effects of nanoparticles on health aren't understood / they could react with things in the body / they could damage cells [1 mark].
- 22.1 Zinc is more reactive than hydrogen [1 mark]. This means zinc forms positive ions more easily than hydrogen [1 mark].
- 22.2 Reduction [1 mark], because the hydrogen ions gain electrons [1 mark].
- 22.3 $4OH^- \rightarrow O_2 + 2H_2O + 4e^-$ [1 mark for correct reactants and products, 1 mark for balancing]
 If you had ' $-4e^-$ ' on the left hand side of the equation instead of ' $+4e^-$ ' on the right, you still get the marks.
- 23.1 Similarity: e.g. they both form positive ions / they both react with acid [1 mark]. Difference: e.g. cobalt has a higher melting point / cobalt forms more than one positive ion / cobalt reacts less vigorously with acid [1 mark].
- 23.2 How to grade your answer:
 Level 0: There is no relevant information. [No marks]
 Level 1: There is a brief description of the similarities and differences between lithium and sodium, but no explanation of these observations. [1 to 2 marks]
 Level 2: There is a detailed comparison of the similarities and differences between lithium and sodium, and some explanation of the observations. [3 to 4 marks]
 Level 3: There is a detailed comparison of the similarities and differences between lithium and sodium, and a good explanation of the observations. [5 to 6 marks]
Here are some points your answer may include:
 Both react to form positive, 1+ ions.
 Both elements are in Group 1, so have one electron in their outer shell.
 Not much energy is needed to remove this one outer electron and give the elements a full outer shell of electrons.
 Both react with acid.
 Sodium reacts more vigorously with acid than lithium.
 Sodium is lower down in the group, so the outer electron in sodium is further away from the nucleus than the outer electron in lithium.
 The attraction between the outer electron and the nucleus of sodium is less than the attraction between the outer electron and the nucleus in lithium.
 Less energy is needed to remove the outer electron of sodium, making it more reactive than lithium.
 Any answer in the range 80–160 °C [1 mark]
- 23.3

